

SHARP ROSENTHAL-TYPE INEQUALITIES FOR MIXTURES AND LOG-CONCAVE VARIABLES

GIORGOS CHASAPIS, ALEXANDROS ESKENAZIS, AND TOMASZ TKOCZ

ABSTRACT. We obtain Rosenthal-type inequalities with sharp constants for moments of sums of independent random variables which are mixtures of a fixed distribution. We also identify extremisers in log-concave settings when the moments of summands are individually constrained.

2020 Mathematics Subject Classification. Primary 60E15; Secondary 60G50, 26D15.

Key words. Rosenthal's inequality, sums of independent random variables, mixtures, log-concave random variables.

1. INTRODUCTION

Rosenthal's inequality, discovered in [35] in connection with questions in the geometry of Banach spaces, provides matching lower and upper bounds for p -norms of sums of independent symmetric random variables in terms of norms of the individual summands, at the expense of a multiplicative constant depending only on p . Recall that a random variable X is symmetric if it has the same distribution as $-X$ (or, equivalently, if it has the same distribution as εX , where ε is a Rademacher variable independent of X). Fix $p > 2$ and let X_1, X_2, \dots be independent symmetric random variables in L_p . Plainly, for any $n \in \mathbb{N}$ we have

$$\left\| \sum_{j=1}^n X_j \right\|_p \geq \left\| \sum_{j=1}^n X_j \right\|_2 = \left(\sum_{j=1}^n \|X_j\|_2^2 \right)^{1/2},$$

where here and throughout $\|Y\|_p = (\mathbb{E}|Y|^p)^{1/p}$ is the p -norm of a random variable Y . Moreover, for independent Rademacher random variables $\varepsilon_1, \varepsilon_2, \dots$ independent of X_1, X_2, \dots , we have

$$\left\| \sum_{j=1}^n X_j \right\|_p^p = \mathbb{E}_X \mathbb{E}_\varepsilon \left| \sum_{j=1}^n \varepsilon_j X_j \right|^p \geq \mathbb{E}_X \left(\mathbb{E}_\varepsilon \left| \sum_{j=1}^n \varepsilon_j X_j \right|^2 \right)^{p/2} = \mathbb{E}_X \left(\sum_{j=1}^n |X_j|^2 \right)^{p/2} \geq \sum_{j=1}^n \mathbb{E}|X_j|^p,$$

thus

$$(1) \quad \left\| \sum_{j=1}^n X_j \right\|_p \geq \max \left\{ \left(\sum_{j=1}^n \|X_j\|_2^2 \right)^{1/2}, \left(\sum_{j=1}^n \|X_j\|_p^p \right)^{1/p} \right\}$$

and the multiplicative constant 1 in front of the maximum in this inequality is clearly optimal (simply consider $n = 1$). Rosenthal in his influential paper [35] established a reversal of (1): for every $2 < p < \infty$, there is a constant C_p which depends only on p such that for every $n \in \mathbb{N}$ and

AE was partially supported by a Junior Research Fellowship from Trinity College, Cambridge. TT's research supported in part by NSF grant DMS-1955175.

every independent symmetric random variables X_1, \dots, X_n in L_p , we have

$$(2) \quad \left\| \sum_{j=1}^n X_j \right\|_p \leq C_p \max \left\{ \left(\sum_{j=1}^n \|X_j\|_2^2 \right)^{1/2}, \left(\sum_{j=1}^n \|X_j\|_p^p \right)^{1/p} \right\}.$$

Rosenthal's original motivation in [35] was to construct new subspaces of $L_p(\mu)$ spaces, answering important questions raised in [25] and [26]. His fundamental inequality (2), which substantially generalises Khinchin's inequality, has also received significant attention in probabilistic literature, prompting several fruitful lines of research. To mention only some in passing, there are generalisations to dependent settings of martingale differences, multilinear forms, and the like (see [4, 13, 17, 6, 30, 32]), random vectors in Banach spaces (see [38, 32]), noncommutative settings (see [20, 19, 21, 22]), as well as many works devoted to optimal inequalities with sharp constants in various setups (see, e.g. [34, 5, 33, 18, 39, 9, 15, 37, 28, 14, 36, 29, 31]).

To describe the last direction in more detail, let \mathbf{C}_p denote the best constant in Rosenthal's inequality, namely the least C_p such that (2) holds for every n and every sequence of independent symmetric random variables X_1, X_2, \dots in L_p . Rosenthal's proof gives $\mathbf{C}_p = O(p^p)$. The sharp behaviour $\mathbf{C}_p = \Theta(p/\log p)$ as $p \rightarrow \infty$ was established by Johnson, Schechtman and Zinn in [18]. This can also be deduced from Latała's precise formula for moments of sums in terms of marginal distributions (see [23, Corollary 3]). The exact value of \mathbf{C}_p is known to be

$$(3) \quad \mathbf{C}_p = \begin{cases} (1 + \|Z\|_p^p)^{1/p}, & 2 < p \leq 4, \\ \left\| \sum_{j=1}^{\xi} \varepsilon_j \right\|_p, & p \geq 4, \end{cases}$$

where Z is a standard Gaussian random variable and ξ is a Poisson random variable with parameter 1, independent of the Rademacher sequence $\varepsilon_1, \varepsilon_2, \dots$. For $p \geq 4$, the value of \mathbf{C}_p was found by Utev in [39] (continuing investigations of Prokhorov from [34] and Pinelis and Utev from [33]). When $2 < p < 4$, (3) was proven by Ibragimov and Sharakhmetov in [15] (building on Utev's approach), and, independently, by Figiel, Hitczenko, Johnson, Schechtman and Zinn in [9] (with different methods treating a more general case of Orlicz functionals in place of moments).

The question of identifying the value of \mathbf{C}_p can be restated as an extremal problem, since

$$(4) \quad \mathbf{C}_p = \sup \left\{ \left\| \sum_{j=1}^n X_j \right\|_p : n \in \mathbb{N} \text{ and } \max \left\{ \sum_{j=1}^n \|X_j\|_2^2, \sum_{j=1}^n \|X_j\|_p^p \right\} \leq 1 \right\}.$$

This reformulation has led to the study of many *Rosenthal-type* extremal problems, in which the supremal value of the p -norm of a sum of independent variables is sought for under more refined assumptions, including constraints of the form $\max \left\{ \sum_{j=1}^n \|X_j\|_2^2, \lambda \sum_{j=1}^n \|X_j\|_p^p \right\} \leq 1$ for suitable parameters $\lambda > 0$, or $\max \{ \|X_j\|_2, \lambda_j \|X_j\|_p \} \leq \mu_j$, where $\lambda_j, \mu_j > 0$. As we shall see in the next section, (asymptotic) maximisers of such multi-constraint problems can frequently be identified which, in turn, often leads to optimal constants in various Rosenthal-type inequalities.

The present paper is concerned with the following question: what is the nature of maximisers of Rosenthal-type extremal problems if we a priori assume that the independent variables X_1, X_2, \dots have additional properties beyond symmetry, such as unimodality or log-concavity? Can one identify the optimal value of the corresponding optimal constant \mathbf{C}_p under these assumptions? Building on Utev's approach from [39], we fully answer these questions in the case of general mixtures and we also characterize the extremising sequences in the log-concave case, using an effective refinement of the one-dimensional localisation principle (see [27, 10, 11]) developed in [8].

2. RESULTS

2.1. General mixtures. Let V be a symmetric random variable. We say that a random variable X is a V -mixture if it has the same distribution as RV for some nonnegative random variable R , independent of V . In particular, when V is a Rademacher random variable, V -mixtures are exactly symmetric random variables and when V is uniform, V -mixtures are exactly symmetric unimodal random variables (see, e.g., [24, Lemma 1]). When V is standard Gaussian random variable, we refer to a V -mixture as a (symmetric) Gaussian mixture (see [2, 7]).

Following Utev [39], for $p > 2$, an integer $n \geq 1$, sequences $a = (a_1, \dots, a_n)$, $b = (b_1, \dots, b_n)$ of positive numbers and positive parameters A, B , we define the classes of n -tuples $X = (X_1, \dots, X_n)$ of independent V -mixtures with constrained moments as follows:

$$(5) \quad \mathcal{M}_V(n, p, a, b) = \left\{ X : \text{each } X_j \text{ is a } V\text{-mixture with } \|X_j\|_2 \leq a_j \text{ and } \|X_j\|_p \leq b_j \right\},$$

and

$$(6) \quad \mathcal{U}_V(n, p, A, B) = \left\{ X : \text{each } X_j \text{ is a } V\text{-mixture and } \sum_{j=1}^n \|X_j\|_2^2 \leq A^2, \sum_{j=1}^n \|X_j\|_p^p \leq B^p \right\},$$

where we have implicitly assumed that $a_j \leq b_j$ for every j (for feasibility). We shall simply write \mathcal{M} and \mathcal{U} to denote \mathcal{M}_V and \mathcal{U}_V with V being a Rademacher random variable, that is for the classes of independent symmetric random variables satisfying the constraints. Moreover, \mathcal{M}'_V and \mathcal{U}'_V will denote the subclasses of \mathcal{M}_V and \mathcal{U}_V respectively, consisting of all the tuples $X = (X_1, \dots, X_n)$ of *identically distributed* V -mixtures satisfying the imposed moment constraints.

Our first theorem is the solution of the following Rosenthal-type extremal problem for mixtures.

Theorem 1. *Fix $A, B > 0$ and let V be a symmetric random variable. For $2 < p < 4$, we have*

$$(7) \quad \sup \left\{ \mathbb{E} \left| \sum_{j=1}^n X_j \right|^p : n \in \mathbb{N} \text{ and } X \in \mathcal{U}_V(n, p, A, B) \right\} = B^p + \|Z\|_p^p A^p,$$

where Z is a standard Gaussian, provided that V is in $L_{p+\delta}$ for some $\delta > 0$. For $p \geq 4$, we have

$$(8) \quad \sup \left\{ \mathbb{E} \left| \sum_{j=1}^n X_j \right|^p : n \in \mathbb{N} \text{ and } X \in \mathcal{U}_V(n, p, A, B) \right\} = \left(\frac{B^p \|V\|_2^2}{A^2 \|V\|_p^p} \right)^{\frac{p}{p-2}} \mathbb{E} \left| \sum_{j=1}^{\xi} V_j \right|^p,$$

provided that V is in L_p , where V_1, V_2, \dots are i.i.d. copies of V and ξ is an independent Poisson random variable with parameter $\left(\frac{A \|V\|_p}{B \|V\|_2} \right)^{\frac{2p}{p-2}} (1 - \mathbb{P}(V = 0))$.

Observe that when $2 < p < 4$, the solution (7) of the extremal problem is independent of the law of V . Theorem 1 readily implies the following optimal Rosenthal inequality for mixtures.

Corollary 2. *The best constant $\mathbf{C}_{p,V}$ such that for any sequence X_1, X_2, \dots of independent V -mixtures and any $n \geq 1$, we have the inequality*

$$\left\| \sum_{j=1}^n X_j \right\|_p \leq \mathbf{C}_{p,V} \max \left\{ \left(\sum_{j=1}^n \|X_j\|_2^2 \right)^{1/2}, \left(\sum_{j=1}^n \|X_j\|_p^p \right)^{1/p} \right\}$$

is

$$(9) \quad \mathbf{C}_{p,V} = \begin{cases} (1 + \|Z\|_p^p)^{1/p}, & 2 < p \leq 4, \\ \left(\frac{\|V\|_2^2}{\|V\|_p^p} \right)^{\frac{1}{p-2}} \left\| \sum_{j=1}^{\xi} V_j \right\|_p, & p \geq 4, \end{cases}$$

where Z is a standard Gaussian random variable, V_1, V_2, \dots are i.i.d. copies of V and ξ is an independent Poisson random variable with parameter $\left(\frac{\|V\|_p}{\|V\|_2}\right)^{\frac{2p}{p-2}} (1 - \mathbb{P}(V = 0))$.

Proof. By its definition and homogeneity, $\mathbf{C}_{p,V}$ satisfies

$$\mathbf{C}_{p,V}^p = \sup \left\{ \mathbb{E} \left| \sum_{j=1}^n X_j \right|^p : n \in \mathbb{N} \text{ and } X \in \mathcal{U}_V(n, p, 1, 1) \right\},$$

and thus (9) follows immediately from (7) and (8) with $A = B = 1$. \square

Theorem 1 specialised to Gaussian mixtures gives the sharp constants in Rosenthal's inequality for sums of nonnegative random variables. This recovers and provides a new proof of the main results obtained independently in [16] by Ibragimov and Sharakhmetov and in [37] by Schechtman which use different approaches: the former adapts Utev's arguments, whereas the latter reduces the problem to sums of independent Poisson random variables using convexity and then optimises over their parameters (in an ingenious way).

Corollary 3 ([16, 37]). *Let $A, B > 0$. We have*

$$(10) \quad \sup \mathbb{E} \left(\sum_{j=1}^n X_j \right)^p = \begin{cases} A^p + B^p, & 1 < p < 2, \\ \left(\frac{B^p}{A}\right)^{\frac{p}{p-1}} \mathbb{E} \xi^p, & p \geq 2, \end{cases}$$

where ξ is a Poisson random variable with parameter $\left(\frac{A}{B}\right)^{\frac{p}{p-1}}$ and the supremum is taken over all $n \geq 1$ and all sequences (X_1, \dots, X_n) of independent, nonnegative random variables with

$$\sum_{j=1}^n \mathbb{E} X_j \leq A \quad \text{and} \quad \sum_{j=1}^n \mathbb{E} X_j^p \leq B^p.$$

Proof. For a fixed sequence of independent positive random variables (X_1, \dots, X_n) , we consider the sequence $(\sqrt{X_1}Z_1, \dots, \sqrt{X_n}Z_n)$ of independent Gaussian mixtures and apply Theorem 1 (we note that $\sum_{j=1}^{\xi} Z_j$ has the same distribution as $\sqrt{\xi}Z_1$). \square

2.2. Log-concave random variables. An important step in Utev's approach leading to the optimal Rosenthal constant (3) for $p > 4$ is a reduction to 3-point distributions: given n and positive sequences a, b , it turns out that $\sup_{X \in \mathcal{M}(n, p, a, b)} \mathbb{E} \left| \sum_{j=1}^n X_j \right|^p$ is attained at the (uniquely determined) 3-point distribution which attains the moment constraints as equalities (for details, see Theorem 8 in the next section). Our main result here, put informally, is a log-concave analogue of this. Recall that a random variable X is log-concave if it has a density of the form $e^{-\phi}$ for a convex function $\phi: \mathbb{R} \rightarrow (-\infty, +\infty]$. Log-concave distributions arise naturally in convex geometry and geometric functional analysis (see e.g. [1, 3]). To state our theorem rigorously, we first need to describe the relevant extremal log-concave distributions (whose convex potentials ϕ turn out to be piecewise linear with at most two pieces on the positive semiaxis).

For two parameters $\alpha \in [0, +\infty)$ and $\gamma \in (0, +\infty]$, we define the log-concave density

$$(11) \quad f_{\alpha, \gamma}(x) = \frac{1}{2(\alpha + 1/\gamma)} \exp \{-\gamma(|x| - \alpha)_+\}$$

with the convention that $\gamma = +\infty$, $\alpha > 0$ gives the uniform density on $[-\alpha, \alpha]$ and that $\alpha = 0$, $\gamma < +\infty$ gives the two-sided exponential density $\frac{\gamma}{2} e^{-\gamma|x|}$. For $\alpha \in (0, +\infty]$ and $\gamma \in [0, +\infty)$, let

$$(12) \quad g_{\alpha, \gamma}(x) = \frac{\gamma}{2(1 - e^{-\alpha\gamma})} \exp \{-\gamma|x|\} \mathbf{1}_{\{|x| \leq \alpha\}}$$

with the convention that $\gamma = 0$, $\alpha > 0$ gives the uniform density on $[-\alpha, \alpha]$ and that $\alpha = +\infty$, $\gamma > 0$ gives the two-sided exponential density $\frac{\gamma}{2}e^{-\gamma|x|}$. We set

$$(13) \quad \mathcal{F}^- = \{f_{\alpha, \gamma} : \alpha \in [0, +\infty), \gamma \in (0, +\infty]\}$$

and

$$(14) \quad \mathcal{F}^+ = \{g_{\alpha, \gamma} : \alpha \in (0, +\infty], \gamma \in [0, +\infty)\}$$

to be the two-parameter family of such densities. The following lemma describes the set of feasible moment parameters. We defer its simple but technical proof to the next section.

Lemma 4. *Let $p > 2$. For every $a, b > 0$ such that*

$$(15) \quad 3^{1/2}(p+1)^{-1/p} \leq \frac{b}{a} \leq 2^{-1/2}\Gamma(p+1)^{1/p},$$

there exist unique $f^- \in \mathcal{F}^-$ and $f^+ \in \mathcal{F}^+$ with

$$(16) \quad \int_{\mathbb{R}} x^2 f^\pm(x) dx = a^2 \quad \text{and} \quad \int_{\mathbb{R}} |x|^p f^\pm(x) dx = b^p.$$

Conversely, the second and p -th moment of any symmetric log-concave variable satisfy (15).

With notation set up, the main theorem of this section is the solution of a Rosenthal-type extremal problem for symmetric log-concave random variables with all moments prescribed.

Theorem 5. *Fix $p > 4$, $n \geq 1$ and let $a_1, \dots, a_n, b_1, \dots, b_n > 0$ be such that each ratio b_j/a_j satisfies (15). Then, we have*

$$(17) \quad \inf \mathbb{E} \left| \sum_{j=1}^n X_j \right|^p = \mathbb{E} \left| \sum_{j=1}^n \mathbf{X}_j^- \right|^p \quad \text{and} \quad \sup \mathbb{E} \left| \sum_{j=1}^n X_j \right|^p = \mathbb{E} \left| \sum_{j=1}^n \mathbf{X}_j^+ \right|^p,$$

where the infimum (respectively supremum) is taken over all sequences $X = (X_1, \dots, X_n)$ of independent symmetric log-concave random variables X_1, \dots, X_n with $\mathbb{E}X_j^2 = a_j^2$ and $\mathbb{E}|X_j|^p = b_j^p$ and the \mathbf{X}_j^- (resp. \mathbf{X}_j^+) have the unique densities from \mathcal{F}^- (resp. \mathcal{F}^+) that satisfy the same constraints.

2.3. Random variables with log-concave tails. A symmetric random variable X is said to have log-concave tails if $T_X(t) = \mathbb{P}(|X| > t)$ is log-concave on $[0, \infty)$. Every log-concave random variable has log-concave tails but the converse is not true, e.g. for Rademacher variables. A modification of the proof of Theorem 5 allows us to also resolve the corresponding Rosenthal-type extremal problems for variables with log-concave tails. Consider the classes of distributions

$$(18) \quad \mathcal{G}^- = \{X : T_X(t) = e^{-a(t-b)_+} \text{ for some } a, b \geq 0\}$$

and

$$(19) \quad \mathcal{G}^+ = \{X : T_X(t) = e^{-at} \mathbf{1}_{[0, b]}(t) \text{ for some } a, b \geq 0\}.$$

Theorem 6. *Fix $p > 4$, $n \geq 1$ and let $a_1, \dots, a_n, b_1, \dots, b_n > 0$ be feasible sequences of second and p -th moments of symmetric random variables with log-concave tails. Then, we have*

$$(20) \quad \inf \mathbb{E} \left| \sum_{j=1}^n X_j \right|^p = \mathbb{E} \left| \sum_{j=1}^n \mathbf{X}_j^- \right|^p \quad \text{and} \quad \sup \mathbb{E} \left| \sum_{j=1}^n X_j \right|^p = \mathbb{E} \left| \sum_{j=1}^n \mathbf{X}_j^+ \right|^p,$$

where the infimum and supremum are taken over all sequences $X = (X_1, \dots, X_n)$ of independent symmetric random variables X_1, \dots, X_n with log-concave tails, $\mathbb{E}X_j^2 = a_j^2$ and $\mathbb{E}|X_j|^p = b_j^p$ and the \mathbf{X}_j^- (resp. \mathbf{X}_j^+) are the unique laws in \mathcal{G}^- (resp. \mathcal{G}^+) that satisfy the same constraints.

3. PROOF OF THEOREM 1

We first quickly show the identity (7) for $2 < p < 4$ as it follows from the case of symmetric random variables (Rademacher mixtures) proven by Utev in [39].

Proof of (7). Since $\mathcal{M}_V \subseteq \mathcal{M}$ and $\mathcal{U}_V \subseteq \mathcal{U}$, thanks to the result for symmetric random variables (see [9, Proposition 8.1] or equation (3) in [15]), we immediately obtain that (7) holds with “ \leq ” in place of “ $=$ ”. To argue that “ \geq ” holds as well, it suffices to modify the example from the proof of [9, Proposition 8.1] by replacing the Rademacher random variables with i.i.d. copies of V . For completeness, we now sketch this construction. Fix $0 < \alpha < A/\|V\|_2$ and set

$$(21) \quad X_j = \begin{cases} \frac{\alpha}{\sqrt{n}}V_j, & 1 \leq j \leq n, \\ \gamma\theta_j V_j, & n < j \leq 2n, \end{cases}$$

where V_1, \dots, V_{2n} are i.i.d. copies of V , $\theta_{n+1}, \dots, \theta_{2n}$ are i.i.d. Bernoulli random variables with parameter λ/n , independent of the V_j , and $\gamma, \lambda > 0$ are parameters to be chosen soon. For $r > 0$,

$$\sum_{j=1}^{2n} \|X_j\|_r^r = \|V\|_r^r (\alpha^r n^{1-r/2} + \gamma^r \lambda).$$

Thus, to ensure that $X \in \mathcal{U}_V(n, p, A, B)$, it suffices to choose γ and λ such that

$$\gamma^2 \lambda = \frac{A^2}{\|V\|_2^2} - \alpha^2 \quad \text{and} \quad \gamma^p \lambda = \frac{B^p}{\|V\|_p^p} - \alpha^p n^{1-p/2}$$

which is clearly possible as long as n is large enough. Using that for arbitrary independent symmetric random variables S, T , we have $\mathbb{E}|S+T|^p \geq \mathbb{E}|S|^p + \mathbb{E}|T|^p$, $p > 2$ (see (1), say), we get

$$\mathbb{E} \left| \sum_{j=1}^{2n} X_j \right|^p \geq \mathbb{E} \left| \sum_{j=1}^n X_j \right|^p + \sum_{j=n+1}^{2n} \mathbb{E}|X_j|^p = \alpha^p \mathbb{E} \left| \frac{\sum_{j=1}^n V_j}{\sqrt{n}} \right|^p + \gamma^p \lambda \|V\|_p^p.$$

The family $\left\{ \left| \frac{\sum_{j=1}^n V_j}{\sqrt{n}} \right|^p \right\}_{n \geq 1}$ is uniformly integrable (since it is bounded in $L_{1+\delta/p}$, by (2), say), so by the central limit theorem, $\mathbb{E} \left| \frac{\sum_{j=1}^n V_j}{\sqrt{n}} \right|^p \rightarrow \|Z\|_p^p \|V\|_2^p$ as $n \rightarrow \infty$. Moreover, $\gamma^p \lambda \|V\|_p^p = B^p - \alpha^p n^{1-p/2} \|V\|_p^p \rightarrow B^p$ as $n \rightarrow \infty$ and letting $\alpha \rightarrow A/\|V\|_2$ finishes the argument. \square

The proof of (8) requires some preparation. We first recall some of Utev’s results. Central to his approach is the following Poissonisation estimate. For a finite nonnegative Borel measure ν on \mathbb{R} , we denote by T_ν a random variable with characteristic function

$$(22) \quad \mathbb{E} e^{itT_\nu} = \exp \left\{ \int_{\mathbb{R}} (e^{itx} - 1) d\nu(x) \right\}, \quad t \in \mathbb{R}.$$

(T_ν can be explicitly constructed as $\sum_{j=1}^\xi X_j$, where X_1, X_2, \dots are i.i.d. copies of a random variable with law $\frac{1}{\nu(\mathbb{R})}\nu$ and ξ is an independent Poisson random variable with parameter $\nu(\mathbb{R})$.) As customary, $\mathcal{B}(\mathbb{R})$ denotes the σ -algebra of all Borel sets in \mathbb{R} .

Theorem 7 (Utev, Theorem 4 in [39]). *Let $\Phi: \mathbb{R} \rightarrow \mathbb{R}$ be an even C^2 function with Φ'' convex. For every $n \in \mathbb{N}$ and independent symmetric random variables X_1, \dots, X_n , we have*

$$(23) \quad \mathbb{E} \Phi \left(\sum_{j=1}^n X_j \right) \leq \mathbb{E} \Phi(T_\nu)$$

with ν is defined by setting

$$(24) \quad \nu(\Gamma) = \sum_{j=1}^n \mathbb{P}(X_j \in \Gamma \setminus \{0\}), \quad \Gamma \in \mathcal{B}(\mathbb{R}).$$

We will also need to use the fact that 3-point distributions are extremal among all symmetric distributions with fixed moments.

Theorem 8 (Utev, Theorem 5 in [39]). *For $p \geq 4$, we have*

$$(25) \quad \sup_{X \in \mathcal{M}(n,p,a,b)} \mathbb{E} \left| \sum_{j=1}^n X_j \right|^p = \mathbb{E} \left| \sum_{j=1}^n \left(\frac{b_j^p}{a_j^2} \right)^{\frac{1}{p-2}} \theta_{j,\mu_j} \varepsilon_j \right|^p,$$

where the θ_{j,μ_j} denote i.i.d. Bernoulli random variables with parameter $\mu_j = (a_j/b_j)^{\frac{2p}{p-2}}$.

We are ready to prove (8). We begin with a lemma showing that i.i.d. sequences are extremal.

Lemma 9. *Let $p \geq 3$, $A, B > 0$ and let V be a symmetric random variable in L_p . We have,*

$$(26) \quad \sup_{n \geq 1, X \in \mathcal{U}_V(n,p,A,B)} \mathbb{E} \left| \sum_{j=1}^n X_j \right|^p = \sup_{n \geq 1, X \in \mathcal{U}'_V(n,p,A,B)} \mathbb{E} \left| \sum_{j=1}^n X_j \right|^p.$$

Proof. A finite nonnegative Borel measure ν on \mathbb{R} is a V -mixture if it is of the form

$$\nu(\Gamma) = \int_0^\infty \mathbb{P}(rV \in \Gamma \setminus \{0\}) d\eta(r), \quad \Gamma \in \mathcal{B}(\mathbb{R})$$

for some finite nonnegative Borel measure η on $(0, \infty)$. We shall argue that both suprema in (26) are equal to the proxy

$$(27) \quad Q = \sup_{\nu \in \mathcal{W}_V(p,A,B)} \mathbb{E}|T_\nu|^p,$$

where

$$\mathcal{W}_V(p, A, B) = \left\{ \nu : \nu \text{ is a } V\text{-mixture with } \nu(\{0\}) = 0, \int_{\mathbb{R}} x^2 d\nu(x) \leq A^2, \int_{\mathbb{R}} |x|^p d\nu(x) \leq B^p \right\}.$$

Step I. We have,

$$\sup_{n \geq 1, X \in \mathcal{U}'_V(n,p,A,B)} \mathbb{E} \left| \sum_{j=1}^n X_j \right|^p \leq \sup_{n \geq 1, X \in \mathcal{U}_V(n,p,A,B)} \mathbb{E} \left| \sum_{j=1}^n X_j \right|^p \leq Q,$$

where the first inequality is clear and the second one follows from (23) for $\Phi(x) = |x|^p$.

Step II. We now have to show that

$$Q \leq \sup_{n \geq 1, X \in \mathcal{U}'_V(n,p,A,B)} \mathbb{E} \left| \sum_{j=1}^n X_j \right|^p.$$

To this end, fix $\nu \in \mathcal{W}_V(p, A, B)$. Take an integer $n \geq \nu(\mathbb{R})$ and i.i.d. random variables X_1, \dots, X_n with the law specified by

$$\mathbb{P}(X_1 \in \Gamma) = \frac{1}{n} \nu(\Gamma), \quad \Gamma \in \mathcal{B}(\mathbb{R}) \text{ with } 0 \notin \Gamma$$

and $\mathbb{P}(X_1 = 0) = 1 - \frac{1}{n} \nu(\mathbb{R})$. Since ν is a V -mixture, the random variables X_j are also i.i.d. V -mixtures. Moreover, $X = (X_1, \dots, X_n) \in \mathcal{U}'_V(n, p, A, B)$, as $\nu \in \mathcal{W}_V(p, A, B)$. Note that the

characteristic function of $\sum_{j=1}^n X_j$ equals

$$\left(1 - \frac{1}{n} \nu(\mathbb{R}) + \frac{1}{n} \int_{\mathbb{R}} e^{itx} d\nu(x)\right)^n = \left(1 + \frac{1}{n} \int_{\mathbb{R}} (e^{itx} - 1) d\nu(x)\right)^n,$$

and thus $\sum_{j=1}^n X_j$ converges in distribution to T_ν as $n \rightarrow \infty$. By Fatou's lemma, we get

$$\mathbb{E}|T_\nu|^p \leq \liminf_{n \rightarrow \infty} \mathbb{E} \left| \sum_{j=1}^n X_j \right|^p$$

which finishes the proof of (26). \square

Proof of (8). By virtue of Lemma 9, we will equivalently show (8) with \mathcal{U}'_V in place of \mathcal{U}_V . First we argue that “ \leq ” holds. To this end, take an n -tuple $X = (X_1, \dots, X_n)$ of i.i.d. V -mixtures in $\mathcal{U}_V(n, p, A, B)$, say $X_j = R_j |V_j| \varepsilon_j$ for some i.i.d. nonnegative random variables R_j and such that the R_j, V_j, ε_j are all independent. Plainly,

$$\mathbb{E} \left| \sum_{j=1}^n X_j \right|^p = \mathbb{E}_V \mathbb{E}_{R, \varepsilon} \left| \sum_{j=1}^n |V_j| R_j \varepsilon_j \right|^p,$$

so conditioning on the values of the V_j and applying (25) yields

$$\begin{aligned} \mathbb{E}_V \mathbb{E}_{R, \varepsilon} \left| \sum_{j=1}^n |V_j| R_j \varepsilon_j \right|^p &\leq \mathbb{E}_V \mathbb{E}_{\theta, \varepsilon} \left| \sum_{j=1}^n \left(\frac{(Bn^{-1/p})^p \|V\|_2^2}{(An^{-1/2})^2 \|V\|_p^p} \right)^{\frac{1}{p-2}} |V_j| \theta_j \varepsilon_j \right|^p \\ &= \left(\frac{B^p \|V\|_2^2}{A^2 \|V\|_p^p} \right)^{\frac{p}{p-2}} \mathbb{E} \left| \sum_{j=1}^n \theta_j V_j \right|^p, \end{aligned}$$

where the θ_j are i.i.d. Bernoulli random variables with parameter

$$\mu = \left(\frac{An^{-1/2} \|V\|_p}{Bn^{-1/p} \|V\|_2} \right)^{\frac{2p}{p-2}} = n^{-1} \left(\frac{A \|V\|_p}{B \|V\|_2} \right)^{\frac{2p}{p-2}}.$$

Let ν be a Borel measure specified by

$$(28) \quad \nu(\Gamma) = \left(\frac{A \|V\|_p}{B \|V\|_2} \right)^{\frac{2p}{p-2}} \mathbb{P}(V \in \Gamma \setminus \{0\}) = n \mathbb{P}(\theta_1 V_1 \in \Gamma \setminus \{0\}), \quad \Gamma \in \mathcal{B}(\mathbb{R}).$$

Then Theorem 7 yields

$$\mathbb{E} \left| \sum_{j=1}^n \theta_j V_j \right|^p \leq \mathbb{E}|T_\nu|^p.$$

Note that T_ν has the same distribution as $\sum_{j=1}^\xi V_j$, where ξ is a Poisson random variable with parameter $\nu(\mathbb{R})$, as in the statement of the theorem. This finishes the proof of the inequality “ \leq ” in (8). The reverse inequality follows by repeating the construction from Step II of the proof of Lemma 9. Alternatively, recalling from the proof of Lemma 9 the notation of the class \mathcal{W}_V as well as the proxy quantity Q , we take the measure ν defined in (28) above and rescale

$$\tilde{\nu}(\Gamma) = \nu \left(\Gamma / \left(\frac{B^p \|V\|_2^2}{A^2 \|V\|_p^p} \right)^{\frac{1}{p-2}} \right), \quad \Gamma \in \mathcal{B}(\mathbb{R})$$

to have, $\tilde{\nu} \in \mathcal{W}_V(p, A, B)$, thus by the definition of Q ,

$$\left(\frac{B^p \|V\|_2^2}{A^2 \|V\|_p^p} \right)^{\frac{p}{p-2}} \mathbb{E}|T_\nu|^p = \mathbb{E}|T_{\tilde{\nu}}|^p \leq Q,$$

which finishes the proof in view of two facts, that T_ν has the same distribution as $\sum_{j=1}^{\xi} V_j$ and that Q equals the supremum from (8) (as shown in the proof of Lemma 9). \square

4. PROOF OF THEOREMS 5 AND 6

4.1. Log-concave random variables. We only consider the case of the infimum and class \mathcal{F}^- , with the obvious modifications left out to address the case of the supremum and \mathcal{F}^+ .

Proof of Lemma 4. Let $f_0, f_1 \in \mathcal{F}^-$ be the uniform and two-sided exponential densities chosen such that $\int_{\mathbb{R}} x^2 f_j(x) dx = a^2$, $j = 0, 1$. Then

$$\int_{\mathbb{R}} |x|^p f_0(x) dx = 3^{p/2} (p+1)^{-1} a^p, \quad \int_{\mathbb{R}} |x|^p f_1(x) dx = 2^{-p/2} \Gamma(p+1) a^p$$

For $\rho \in [0, +\infty]$, let

$$\gamma = \gamma(\rho) = a^{-1} \sqrt{2 + \frac{\rho^3 + 3\rho^2}{3(\rho+1)}}$$

which is chosen such that the density $g_\rho(x) = f_{\frac{a}{\gamma}, \gamma}$ satisfies

$$\int_{\mathbb{R}} x^2 g_\rho(x) dx = a^2.$$

Since $g_0 = f_1$ and $g_\infty = f_0$, the intermediate value property gives the existence of ρ such that $\int_{\mathbb{R}} |x|^p g_\rho(x) dx = b^p$, as desired. The uniqueness follows from the fact that two arbitrary distinct densities from \mathcal{F}^- intersect each other at most twice on $(0, \infty)$. The converse implication is classical, see for instance [8, Remark 14] and [12, Proposition 5.5]. \square

We need several more ancillary results.

Lemma 10. *Let g be an even log-concave density on \mathbb{R} and let $f \in \mathcal{F}^-$. Then $g - f$ changes sign at most 3 times on $(0, +\infty)$.*

Proof. Say $f = f_{\alpha, \gamma}$. Examining $\log g - \log f$ on $(0, +\infty)$, this difference clearly changes sign at most once on $(0, \alpha]$ (by monotonicity of $\log g$, since $\log f$ is constant there) and at most twice on $[\alpha, +\infty)$ (by concavity of $\log g$, since $\log f$ is linear there). \square

Lemma 11. *Let $p > 4$. The following function*

$$(29) \quad \psi_p(x) = |\sqrt{x} + 1|^p + |\sqrt{x} - 1|^p - 2x^{p/2}, \quad x \geq 0$$

is strictly convex on $(0, +\infty)$.

Proof. Let $g(x) = |x+1|^p + |x-1|^p - 2x^{p/2}$, $x \geq 0$. We have $\psi_p(x) = g(\sqrt{x})$ and $\psi'_p(x) = \frac{g'(\sqrt{x})}{2\sqrt{x}}$. Since $g'(0) = 0$, it suffices to show that g' is convex, because then ψ'_p is increasing. For $x > 1$, we have

$$\frac{1}{p(p-1)(p-2)} g'''(x) = (x+1)^{p-3} + (x-1)^{p-3} - 2x^{p-3} > 0,$$

by convexity. For $0 < x < 1$, we have

$$\begin{aligned} \frac{1}{p(p-1)(p-2)(p-3)} g'''(x) &= \frac{(x+1)^{p-3} - (1-x)^{p-3} - 2x^{p-3}}{p-3} \\ &= \int_{1-x}^{1+x} t^{p-4} dt - 2 \int_0^x t^{p-4} dt > 0, \end{aligned}$$

by the monotonicity of the integrand. \square

Lemma 12. For $0 < x_1 < x_2 < x_3$ and a convex function $\phi: (0, +\infty) \rightarrow \mathbb{R}$, we have

$$(30) \quad \det \begin{bmatrix} 1 & x_1 & \phi(x_1) \\ 1 & x_2 & \phi(x_2) \\ 1 & x_3 & \phi(x_3) \end{bmatrix} \geq 0.$$

Moreover, the inequality is strict if ϕ is strictly convex.

Proof. The desired inequality is equivalent to

$$\frac{x_3 - x_2}{x_3 - x_1} \phi(x_1) + \frac{x_2 - x_1}{x_3 - x_1} \phi(x_3) \geq \phi(x_2)$$

which clearly follows from convexity. \square

Lemma 13. Let $p > 4$, let α, β, γ be real numbers. The function

$$h(x) = |x + 1|^p + |x - 1|^p - \alpha - \beta x^2 - \gamma x^p$$

has at most 3 sign changes on $(0, +\infty)$. Moreover, if it has exactly 3, then its signature is $+, -, +, -$.

Proof. It suffices to consider the case when h has 3 distinct sign changes, say at $0 < x_1 < x_2 < x_3 < \infty$ and show that it does not have any more. First note that then, under the notation used in Lemma 11,

$$\gamma - 2 = \frac{\det \begin{bmatrix} 1 & x_1^2 & |x_1+1|^p + |x_1-1|^p \\ 1 & x_2^2 & |x_2+1|^p + |x_2-1|^p \\ 1 & x_3^2 & |x_3+1|^p + |x_3-1|^p \end{bmatrix}}{\det \begin{bmatrix} 1 & x_1^2 & x_1^p \\ 1 & x_2^2 & x_2^p \\ 1 & x_3^2 & x_3^p \end{bmatrix}} - 2 = \frac{\det \begin{bmatrix} 1 & x_1^2 & \psi_p(x_1^2) \\ 1 & x_2^2 & \psi_p(x_2^2) \\ 1 & x_3^2 & \psi_p(x_3^2) \end{bmatrix}}{\det \begin{bmatrix} 1 & x_1^2 & x_1^p \\ 1 & x_2^2 & x_2^p \\ 1 & x_3^2 & x_3^p \end{bmatrix}},$$

where we used the linearity of the determinant with respect to the last column. It follows from Lemmas 11 and 12 that both the numerator and the denominator are positive and thus $\gamma > 2$. In particular, this gives that $h^{(k)}(+\infty) = -\infty$, for $k = 0, 1, 2, 3$. Furthermore,

$$h'''(x) = p(p-1)(p-2)x^{p-3}(|1+x^{-1}|^{p-3} + |1-x^{-1}|^{p-3} \operatorname{sgn}(1-x^{-1}) - \gamma)$$

and we verify that the function in the brackets changes sign exactly once on $(0, +\infty)$ from $+$ to $-$. As $h''(+\infty) = -\infty$, we gather that h'' changes sign either only once on $(0, +\infty)$, in which case the conclusion follows, or exactly twice, with the signature $-, +, -$. In the latter case, since $h'(0) = 0$ and $h'(+\infty) = -\infty$, we get at most two sign changes of h' on $(0, +\infty)$, hence h has at most 3 sign changes on $(0, +\infty)$. Since $h(+\infty) = -\infty$, the signature of h is $+, -, +, -$. \square

Proof of Theorem 5. Fix a sequence $X = (X_1, \dots, X_n)$ as in the statement, an index $1 \leq j \leq n$ and a parameter $z \in \mathbb{R}$. Thanks to independence, it suffices to show that

$$(31) \quad \mathbb{E}|X_j + z|^p \geq \mathbb{E}|\mathbf{X}_j^- + z|^p.$$

Indeed, if (31) holds, then swapping one variable at a time, we get

$$\mathbb{E} \left| \sum_{j=1}^n X_j \right|^p \geq \mathbb{E} \left| \mathbf{X}_1^- + \sum_{j=2}^n X_j \right|^p \geq \dots \geq \mathbb{E} \left| \sum_{j=1}^n \mathbf{X}_j^- \right|^p,$$

which is exactly (17). To prove (31), let f_j be the density of X_j and \mathbf{f}_j^- be the density of \mathbf{X}_j^- . Using symmetry, the inequality in question is equivalent to

$$\int_0^\infty (f_j(x) - \mathbf{f}_j^-(x)) \cdot \phi(x) \, dx \geq 0,$$

where $\phi(x) = |x + z|^p + |x - z|^p$. To prove it, we employ the technique of “interlacing densities”. Since the integrals of f_j and \mathbf{f}_j^- against 1 , x^2 and x^p are the same, for every $\alpha, \beta, \gamma \in \mathbb{R}$, the inequality above is equivalent to

$$\int_0^\infty (f_j(x) - \mathbf{f}_j^-(x)) \cdot (\phi(x) - \alpha - \beta x^2 - \gamma x^p) dx \geq 0.$$

For the same reason, $f_j - \mathbf{f}_j^-$ need change sign at least 3 times on $(0, +\infty)$. In view of Lemma 10, there are exactly 3 sign changes, say at $x_1 < x_2 < x_3$ and the signature must necessarily be $+, -, +, -$ (because the first sign change will occur on the interval where \mathbf{f}_j^- is constant and f_j decreases on $(0, +\infty)$). We choose α, β, γ such that $f(x) = \phi(x) - \alpha - \beta x^2 - \gamma x^p$ vanishes at x_1, x_2, x_3 . By Lemma 13, h in fact changes sign on $(0, +\infty)$ exactly at those roots and has signature $+, -, +, -$. Thus the integrand is pointwise nonnegative and (31) follows. \square

4.2. Log-concave tails. The proof of Theorem 5 with minor modifications also gives Theorem 6. Skipping most of the details, the main point is that now we write

$$\begin{aligned} \mathbb{E}|X_j + z|^p - \mathbb{E}|\mathbf{X}_j^- + z|^p &= \int_0^\infty \phi'(t) (T_{X_j}(t) - T_{\mathbf{X}_j^-}(t)) dt \\ &= \int_0^\infty (\phi'(t) - \beta t - \gamma t^{p-1}) \cdot (T_{X_j}(t) - T_{\mathbf{X}_j^-}(t)) dt, \end{aligned}$$

where the second equality uses the constraints on the second and p th moments. Since $\mathbf{X}_j^- \in \mathcal{G}^-$ (see also the definition of the classes \mathcal{L}_2^\pm in [8]), the difference $T_{X_j} - T_{\mathbf{X}_j^-}$ changes sign exactly 2 times on $(0, +\infty)$ with signature $-, +, -$. On the other hand, choosing β and γ such that $\phi'(t) - \beta t - \gamma t^{p-1}$ vanishes at those sign change points, the proof of Lemma 13 yields that this bracket has exactly these sign changes with signature $-, +, -$ and we arrive at the conclusion of Theorem 6. To get the reverse inequality (the supremum instead of the infimum), we choose $\mathbf{X}_j^+ \in \mathcal{G}^+$ and repeat the argument while noting the different sign patterns. \square

5. FURTHER REMARKS AND DIRECTIONS

5.1. More constraints. With the aid of results from [8], the arguments of the previous section for log-concave tails can be extended to handle more than two moment constraints, that is given n and say ℓ constraints and feasible $a_{j,k} > 0$, $j = 1, \dots, n$, $k = 1, \dots, \ell$, the infimum (respectively supremum) is taken over all sequences (X_1, \dots, X_n) of length n consisting of independent symmetric random variables with log-concave tails satisfying $\mathbb{E}|X_j|^{p_1} = a_{j,1}, \dots, \mathbb{E}|X_j|^{p_\ell} = a_{j,\ell}$, $j = 1, \dots, n$. The set of feasible parameters $a_{j,k}$ is described in precisely [8, Theorem 7]. We believe that with some extra work, the same extension is possible in the setting of Theorem 5, that is for symmetric log-concave random variables.

5.2. Sharp constants. It has been elusive to us how to obtain an analogue of Theorem 1 for symmetric log-concave random variables. We find it an interesting and challenging problem. The natural approach, via Theorem 5, leaves us with an optimisation problem over the a_j and b_j , unclear how to tackle.

REFERENCES

- [1] Shiri Artstein-Avidan, Apostolos Giannopoulos, and Vitali D. Milman. *Asymptotic geometric analysis. Part I*, volume 202 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2015.
- [2] Roland Averkamp and Christian Houdré. Wavelet thresholding for non-necessarily Gaussian noise: idealism. *Ann. Statist.*, 31(1):110–151, 2003.

- [3] Silouanos Brazitikos, Apostolos Giannopoulos, Petros Valettas, and Beatrice-Helen Vritsiou. *Geometry of isotropic convex bodies*, volume 196 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2014.
- [4] Donald L. Burkholder. Distribution function inequalities for martingales. *Ann. Probability*, 1:19–42, 1973.
- [5] David C. Cox and Johannes H. B. Kemperman. Sharp bounds on the absolute moments of a sum of two i.i.d. random variables. *Ann. Probab.*, 11(3):765–771, 1983.
- [6] Victor H. de la Peña, Rustam Ibragimov, and Shaturgun Sharakhmetov. On extremal distributions and sharp L_p -bounds for sums of multilinear forms. *Ann. Probab.*, 31(2):630–675, 2003.
- [7] Alexandros Eskenazis, Piotr Nayar, and Tomasz Tkocz. Gaussian mixtures: entropy and geometric inequalities. *Ann. Probab.*, 46(5):2908–2945, 2018.
- [8] Alexandros Eskenazis, Piotr Nayar, and Tomasz Tkocz. Sharp comparison of moments and the log-concave moment problem. *Adv. Math.*, 334:389–416, 2018.
- [9] Tadeusz Figiel, Paweł Hitczenko, William B. Johnson, Gideon Schechtman, and Joel Zinn. Extremal properties of Rademacher functions with applications to the Khintchine and Rosenthal inequalities. *Trans. Amer. Math. Soc.*, 349(3):997–1027, 1997.
- [10] Matthieu Fradelizi and Olivier Guédon. The extreme points of subsets of s -concave probabilities and a geometric localization theorem. *Discrete Comput. Geom.*, 31(2):327–335, 2004.
- [11] Matthieu Fradelizi and Olivier Guédon. A generalized localization theorem and geometric inequalities for convex bodies. *Adv. Math.*, 204(2):509–529, 2006.
- [12] Olivier Guédon, Piotr Nayar, and Tomasz Tkocz. Concentration inequalities and geometry of convex bodies. In *Analytical and probabilistic methods in the geometry of convex bodies*, volume 2 of *IMPAN Lect. Notes*, pages 9–86. Polish Acad. Sci. Inst. Math., Warsaw, 2014.
- [13] Paweł Hitczenko. Best constants in martingale version of Rosenthal’s inequality. *Ann. Probab.*, 18(4):1656–1668, 1990.
- [14] Marat Ibragimov and Rustam Ibragimov. Optimal constants in the Rosenthal inequality for random variables with zero odd moments. *Statist. Probab. Lett.*, 78(2):186–189, 2008.
- [15] Rustam Ibragimov and Shaturgun Sharakhmetov. On an exact constant for the Rosenthal inequality. *Teor. Veroyatnost. i Primenen.*, 42(2):341–350, 1997.
- [16] Rustam Ibragimov and Shaturgun Sharakhmetov. The best constant in the Rosenthal inequality for nonnegative random variables. *Statist. Probab. Lett.*, 55(4):367–376, 2001.
- [17] Rustam Ibragimov, Shaturgun Sharakhmetov, and Aydin Cecen. Exact estimates for moments of random bilinear forms. *J. Theoret. Probab.*, 14(1):21–37, 2001.
- [18] William B. Johnson, Gideon Schechtman, and Joel Zinn. Best constants in moment inequalities for linear combinations of independent and exchangeable random variables. *Ann. Probab.*, 13(1):234–253, 1985.
- [19] Marius Junge, Javier Parcet, and Quanhua Xu. Rosenthal type inequalities for free chaos. *Ann. Probab.*, 35(4):1374–1437, 2007.
- [20] Marius Junge and Quanhua Xu. Noncommutative Burkholder/Rosenthal inequalities. *Ann. Probab.*, 31(2):948–995, 2003.
- [21] Marius Junge and Quanhua Xu. Noncommutative Burkholder/Rosenthal inequalities. II. Applications. *Israel J. Math.*, 167:227–282, 2008.
- [22] Marius Junge and Qiang Zeng. Noncommutative Bennett and Rosenthal inequalities. *Ann. Probab.*, 41(6):4287–4316, 2013.
- [23] Rafał Łatała. Estimation of moments of sums of independent real random variables. *Ann. Probab.*, 25(3):1502–1513, 1997.
- [24] Rafał Łatała and Krzysztof Oleszkiewicz. A note on sums of independent uniformly distributed random variables. *Colloq. Math.*, 68(2):197–206, 1995.
- [25] Joram Lindenstrauss and Aleksander Pełczyński. Absolutely summing operators in L_p -spaces and their applications. *Studia Math.*, 29:275–326, 1968.
- [26] Joram Lindenstrauss and Haskell P. Rosenthal. The \mathcal{L}_p spaces. *Israel J. Math.*, 7:325–349, 1969.
- [27] László Lovász and Miklós Simonovits. Random walks in a convex body and an improved volume algorithm. *Random Structures Algorithms*, 4(4):359–412, 1993.
- [28] Iosif Pinelis. Exact inequalities for sums of asymmetric random variables, with applications. *Probab. Theory Related Fields*, 139(3-4):605–635, 2007.

- [29] Iosif Pinelis. Exact Rosenthal-type inequalities for $p = 3$, and related results. *Statist. Probab. Lett.*, 83(12):2634–2637, 2013.
- [30] Iosif Pinelis. Best possible bounds of the von Bahr–Esseen type. *Ann. Funct. Anal.*, 6(4):1–29, 2015.
- [31] Iosif Pinelis. Exact Rosenthal-type bounds. *Ann. Probab.*, 43(5):2511–2544, 2015.
- [32] Iosif Pinelis. Rosenthal-type inequalities for martingales in 2-smooth Banach spaces. *Theory Probab. Appl.*, 59(4):699–706, 2015.
- [33] Iosif F. Pinelis and Sergey A. Utev. Estimates of moments of sums of independent random variables. *Teor. Veroyatnost. i Primenen.*, 29(3):554–557, 1984.
- [34] Yuriĭ V. Prokhorov. An extremal problem in probability theory. *Theor. Probability Appl.*, 4:201–203, 1959.
- [35] Haskell P. Rosenthal. On the subspaces of L^p ($p > 2$) spanned by sequences of independent random variables. *Israel J. Math.*, 8:273–303, 1970.
- [36] Pavel S. Ruzankin. On Cox-Kemperman moment inequalities for independent centered random variables. *Statist. Probab. Lett.*, 86:80–84, 2014.
- [37] Gideon Schechtman. Extremal configurations for moments of sums of independent positive random variables. In *Banach spaces and their applications in analysis*, pages 183–191. Walter de Gruyter, Berlin, 2007.
- [38] Michel Talagrand. Isoperimetry and integrability of the sum of independent Banach-space valued random variables. *Ann. Probab.*, 17(4):1546–1570, 1989.
- [39] Sergey A. Utev. Extremal problems in moment inequalities. In *Limit theorems of probability theory*, volume 5 of *Trudy Inst. Mat.*, pages 56–75, 175. “Nauka” Sibirsk. Otdel., Novosibirsk, 1985.

(G. C.) AND (T. T.) CARNEGIE MELLON UNIVERSITY; PITTSBURGH, PA 15213, USA.

Email address: {gchasapi,ttkocz}@andrew.cmu.edu

(A. E.) CNRS, INSTITUT DE MATHÉMATIQUES DE JUSSIEU, SORBONNE UNIVERSITÉ, FRANCE AND TRINITY COLLEGE, UNIVERSITY OF CAMBRIDGE, UK.

Email address: alexandros.eskenazis@imj-prg.fr, ae466@cam.ac.uk