

# IMPROVED BOUNDS FOR HADWIGER'S COVERING PROBLEM VIA THIN-SHELL ESTIMATES

HAN HUANG, BOAZ A. SLOMKA, TOMASZ TKOCZ, AND BEATRICE-HELEN VRITSIOU

ABSTRACT. A central problem in discrete geometry, known as Hadwiger's covering problem, asks what the smallest natural number  $N(n)$  is such that every convex body in  $\mathbb{R}^n$  can be covered by a union of the interiors of at most  $N(n)$  of its translates. Despite continuous efforts, the best general upper bound known for this number remains as it was more than sixty years ago, of the order of  $\binom{2n}{n}n \ln n$ .

In this note, we improve this bound by a sub-exponential factor. That is, we prove a bound of the order of  $\binom{2n}{n}e^{-c\sqrt{n}}$  for some universal constant  $c > 0$ .

Our approach combines ideas from [3] by Artstein-Avidan and the second named author with tools from Asymptotic Geometric Analysis. One of the key steps is proving a new lower bound for the maximum volume of the intersection of a convex body  $K$  with a translate of  $-K$ ; in fact, we get the same lower bound for the volume of the intersection of  $K$  and  $-K$  when they both have barycenter at the origin. To do so, we make use of measure concentration, and in particular of thin-shell estimates for isotropic log-concave measures.

Using the same ideas, we establish an exponentially better bound for  $N(n)$  when restricting our attention to convex bodies that are  $\psi_2$ . By a slightly different approach, an exponential improvement is established also for classes of convex bodies with positive modulus of convexity.

## 1. INTRODUCTION

**1.1. Hadwiger's covering problem.** A long-standing problem in discrete geometry asks whether every convex body in  $\mathbb{R}^n$  can be covered by a union of at most  $2^n$  translates of its interior. It also asks whether  $2^n$  translates are needed only for affine images of the  $n$ -cube.

This problem was posed by Hadwiger [13] for  $n \geq 3$  but was already considered and settled for  $n = 2$  a few years earlier by Levi [16]. An equivalent formulation, in which the interior of the convex body is replaced by smaller homothetic copies of it, was independently posed by Gohberg and Markus [10]. Other equivalent formulations of this problem were posed by Hadwiger [14] and Boltyanski [6] in terms of illuminating the boundary of a convex body by outer light sources. For a comprehensive survey of this problem and most of the progress made so far towards its solution see e.g. [7, 4, 23].

Putting things formally, a subset of  $\mathbb{R}^n$  is called a convex body if it is a compact convex set with non-empty interior. The covering number of a set  $A \subseteq \mathbb{R}^n$  by a set

$B \subseteq \mathbb{R}^n$  is given by

$$N(A, B) = \min \left\{ N \in \mathbb{N} : \exists x_1, \dots, x_N \in \mathbb{R}^n \text{ such that } A \subseteq \bigcup_{i=1}^N \{x_i + B\} \right\},$$

where  $x + B = \{x + b : b \in B\}$ . Denoting the interior of  $B$  by  $\text{int } B$  and letting  $\lambda B = \{\lambda b : b \in B\}$  for  $\lambda \in \mathbb{R}$ , Hadwiger's conjecture states the following.

**Conjecture.** *Let  $K \subseteq \mathbb{R}^n$  be a convex body. Then for some  $0 < \lambda < 1$  one has  $N(K, \lambda K) \leq 2^n$ , or equivalently  $N(K, \text{int } K) \leq 2^n$ . Moreover, equality holds only if  $K$  is an affine image of the  $n$ -cube.*

The currently best general upper bound known for  $n \geq 3$  is  $\binom{2n}{n}(n \ln n + n \ln \ln n + 5n)$ , while the best bound for centrally-symmetric convex bodies (i.e. convex bodies  $K$  satisfying  $K = -K$ ) is  $2^n(n \ln n + n \ln \ln n + 5n)$ . Both bounds are simple consequences of Rogers' estimates [25] for the asymptotic lower density of a covering of the whole space by translates of a general convex body, combined with the Rogers-Shephard inequality [26], as can be seen in [8] and [27]. For small  $n$ , there are better bounds too (see e.g. [24]).

A fractional version of the illumination problem was considered by Naszódi [21], where the upper bounds of  $2^n$  for the centrally-symmetric case, and  $\binom{2n}{n}$  for the general case were obtained. The same bounds, as well as the extremity of the  $n$ -cube in the centrally-symmetric case, were established by Artstein-Avidan and the second named author in [3] by considering fractional covering numbers of convex bodies. Moreover, together with an inequality linking integral covering numbers and fractional covering numbers (see Section 3 below), the aforementioned best known upper bounds for Hadwiger's classical problem were recovered (technically, only the bound in the centrally-symmetric case was explicitly recovered, but the proof of the general bound is almost verbatim the same). These bounds were recovered once more in [17]. For additional recent results on Hadwiger's problem, see [29], [18], and references therein.

**1.1.1. Main results.** We combine ideas from [3] with a new result on the Kövner-Besicovitch measure of symmetry for convex bodies, which we discuss in Section 1.2. As a result, we obtain a new general upper bound for Hadwiger's problem:

**Theorem 1.1.** *There exist universal constants  $c_1, c_2 > 0$  such that for all  $n \geq 2$  and every convex body  $K \subseteq \mathbb{R}^n$ , one has*

$$N(K, \text{int } K) \leq c_1 4^n e^{-c_2 \sqrt{n}}.$$

For  $\psi_2$  bodies (for definitions and more details see Section 2 below), we obtain the following exponential improvement:

**Theorem 1.2.** *Let  $K \subseteq \mathbb{R}^n$  be a convex body with barycenter at the origin which is  $\psi_2$  with constant  $b_2 > 0$ . Then*

$$N(K, \text{int}K) \leq c_1 4^n e^{-c_2 b_2^{-2} n}.$$

**1.2. The Kövner-Besicovitch measure of symmetry.** Denote the set of all convex bodies in  $\mathbb{R}^n$  by  $\mathcal{K}_n$ . Denote the Lebesgue volume of a measurable set  $A \subseteq \mathbb{R}^n$  by  $|A|$ .

Let  $K \subseteq \mathbb{R}^n$  be a convex body. Given a point  $x \in \mathbb{R}^n$ , let us call here the set  $(K - x) \cap (x - K)$  the *symmetric intersection* of  $K$  at  $x$ . As defined by Grünbaum [11], the following is a measure of symmetry for  $K$ , referred to as the Kövner-Besicovitch measure of symmetry:

$$\Delta_{KB}(K) = \max_{x \in \mathbb{R}^n} \frac{|(K - x) \cap (x - K)|}{|K|} = \max_{x \in \mathbb{R}^n} \frac{|K \cap (x - K)|}{|K|}.$$

To study this quantity, throughout this paper, we use the fact that the volume of the symmetric intersection of a convex body at a point  $x$  is the same as its convolution square at  $2x$ , i.e., the convolution relation

$$|(K - x) \cap (x - K)| = |K \cap (2x - K)| = (\mathbb{1}_K * \mathbb{1}_K)(2x),$$

where  $\mathbb{1}_K$  is the indicator function of  $K$ . Combining this with the fact that the support of  $\mathbb{1}_K * \mathbb{1}_K$  is  $2K$ , one easily obtains by integration that

$$(1.1) \quad \min_{K \in \mathcal{K}_n} \Delta_{KB}(K) \geq 2^{-n}.$$

Denote by  $b(K)$  the barycenter of  $K$ . By fixing this as the point of reference, one may consider the volume ratio of the symmetric intersection of  $K$  at its barycenter as another measure of symmetry for  $K$ . A result of V. Milman and Pajor [20] tells us that

$$(1.2) \quad \frac{|(K - b(K)) \cap (b(K) - K)|}{|K|} \geq 2^{-n}.$$

The optimal lower bound, in both instances, is not known and conjectured to be attained by the simplex, which would imply a lower bound of the order of  $(\frac{2}{e})^n$  (see e.g. [11], [28] for more details).

**1.2.1. A new lower bound.** Our second goal in this note is to improve both (1.1) and (1.2). We consider two approaches, both of which involve using the property of a (properly normalized) log-concave measure to concentrate in a thin-shell, and in particular a quantitative form of it by Guédon and E. Milman [12]. More precisely, let  $X$  and  $Y$  be independent random vectors, uniformly distributed on a convex body  $K \subseteq \mathbb{R}^n$ . Our first approach is based on the comparison of the measure of a ball, whose boundary is between the two thin shells around which the distributions

of  $X$  and  $(X + Y)/2$  are concentrated, according to each of these measures; this leads to the improvement of (1.1).

The second approach, which allows us to bound the volume of the symmetric intersection of  $K$  at its barycenter and to improve (1.2), combines the above mentioned thin-shell estimates of Guédon and E. Milman with the notion of entropy. Given that there is not much reason to believe our bounds are optimal, we have chosen to present both approaches since either might have the potential to give further improvements.

To turn to details, we prove the following:

**Theorem 1.3.** *For some universal constant  $c > 0$ , we have*

$$\min_{K \in \mathcal{K}_n} \Delta_{KB}(K) \geq \min_{K \in \mathcal{K}_n : b(K)=0} \frac{|K \cap (-K)|}{|K|} \geq \frac{\exp(cn^{1/2})}{2^n}.$$

Theorem 1.3 is a particular consequence of Propositions 2.2 and 5.3 below, which provide a lower bound for  $\Delta_{KB}(K)$  and  $|K \cap (-K)|/|K|$  by taking into account the  $\psi_\alpha$  behavior of the convex body  $K$  (for definitions and more details see Section 2 below). In particular, for  $\psi_2$  bodies, we have the following exponential improvement of (1.1) and 1.2.

**Corollary 1.4.** *(of Propositions 2.2 and 5.3) Let  $K \in \mathbb{R}^n$  be a convex body centered at the origin which is  $\psi_2$  with constant  $b_2 > 0$ . Then*

$$\Delta_{KB}(K) \geq \frac{|K \cap (-K)|}{|K|} \geq \frac{\exp(cb_2^{-2}n)}{2^n}.$$

**1.3. Positive modulus of convexity.** The modulus of convexity of a centered convex body  $K \subseteq \mathbb{R}^n$  is defined by

$$\delta_K(\varepsilon) = \inf \left\{ 1 - \left\| \frac{x+y}{2} \right\|_K : \|x\|_K, \|y\|_K \leq 1, \|x-y\|_K \geq \varepsilon \right\},$$

where  $\|x\|_K = \inf\{r > 0 : x \in rK\}$  is the gauge function of  $K$ . We say that  $K$  is uniformly convex if  $\delta_K(\varepsilon) > 0$  for all  $0 < \varepsilon < 2$ . Note that in the finite-dimensional case,  $K \subseteq \mathbb{R}^n$  is strictly convex (i.e. the boundary of  $K$  contains no line segments) if and only if it is uniformly convex.

Using a different concentration result of Arias-De-Reyna, Ball, and Villa [1], which was generalized by Gluskin and Milman [9], we extend Theorems 1.2 and 1.4 to the class of convex bodies whose modulus of convexity is positive for some  $0 < \varepsilon < \sqrt{2}$ . More precisely, for  $0 < r < 1$  and  $0 < \varepsilon < \sqrt{2}$ , let  $\mathcal{K}_{n,r,\varepsilon}$  be the class of centered convex bodies  $K \subseteq \mathbb{R}^n$  for which  $\delta_K(\varepsilon) \geq r$ .

**Theorem 1.5.** *Let  $0 < r < 1$ ,  $0 < \varepsilon < \sqrt{2}$ , and let  $K \in \mathcal{K}_{n,r,\varepsilon}$ . Then, for  $\alpha := 1 - \exp\left(-\frac{(\sqrt{2}-\varepsilon)^2}{4}n\right)$ , we have*

$$\Delta_{KB}(K) \geq \alpha 2^{-n} \left( \frac{1}{1-r} \right)^n, \quad \frac{|K \cap (-K)|}{|K|} \geq \frac{1}{e\sqrt{n}} 2^{-n} \left( \frac{1}{1-\alpha r} \right)^n.$$

**Theorem 1.6.** *Let  $0 < r < 1$ ,  $0 < \varepsilon < \sqrt{2}$ , and let  $K \in \mathcal{K}_{n,r,\varepsilon}$ . Then*

$$N(K, \text{int}K) \leq \left( 1 - e^{-\frac{(\sqrt{2}-\varepsilon)^2}{4}n} \right)^{-1} (4(1-r))^n.$$

The paper is organized as follows. In Section 2 we prove the first part of Theorem 1.3 and of Corollary 1.4 (the bounds for the Kövner-Besicovitch measure of symmetry), and in Section 3 we apply these to Hadwiger's covering problem. Section 4 is devoted to the respective bounds in the case of uniformly convex bodies, i.e. the first part of Theorem 1.5 as well as Theorem 1.6. Finally, in Section 5 we complete the proofs of Theorems 1.3 and 1.5 and of Corollary 1.4 by showing via our second approach how to bound the volume of the symmetric intersection of  $K$  at its barycenter as well. A couple of concluding remarks are gathered at the end.

## 2. BOUNDING THE CONVOLUTION SQUARE

This section is devoted to the proof of Proposition 2.2 below. To that end, we need to recall some facts and results.

Denote the standard Euclidean inner product on  $\mathbb{R}^n$  by  $\langle \cdot, \cdot \rangle$ , and the corresponding Euclidean norm on  $\mathbb{R}^n$  by  $\|\cdot\|_2$ . We shall also denote probability by  $\mathbb{P}$  and expectation by  $\mathbb{E}$ .

Recall that a random vector in  $\mathbb{R}^n$  is called isotropic if  $\mathbb{E}X = 0$  (i.e., its barycenter is the origin) and  $\mathbb{E}(X \otimes X) = Id$  (i.e., its covariance matrix is the identity). We say that  $X$  is  $\psi_\alpha$  with constant  $b_\alpha$  if

$$(\mathbb{E}|\langle X, y \rangle|^p)^{1/p} \leq b_\alpha p^{1/\alpha} (\mathbb{E}|\langle X, y \rangle|^2)^{1/2} \quad \forall p \geq 2, \quad \forall y \in \mathbb{R}^n.$$

A function  $f : \mathbb{R}^n \rightarrow [0, \infty)$  is called log-concave if  $\log f$  is concave on the support of  $f$ . It is well-known that any random vector  $X$  in  $\mathbb{R}^n$  with a log-concave density is  $\psi_1$  with  $b_1 \leq C$ , for some universal constant  $C > 0$  (see e.g. [2, p. 115]).

We shall need the following thin-shell deviation estimate of Guédon and E. Milman:

**Theorem 2.1.** ([12, Theorem 1.1]) *Let  $X$  denote an isotropic random vector in  $\mathbb{R}^n$  with log-concave density, which is in addition  $\psi_\alpha$  ( $\alpha \in [1, 2]$ ) with constant  $b_\alpha$ . Then,*

$$\mathbb{P}(\|X\|_2 - \sqrt{n} \geq t\sqrt{n}) \leq C \exp(-c'b_\alpha^{-\alpha} \min(t^{2+\alpha}, t)n^{\alpha/2}) \quad \forall t \geq 0,$$

where  $c' > 0$  is some universal constant.

We remark that the dependence in  $n$  in Theorem 2.1 is optimal, while the dependence in  $t$  was recently improved by Lee and Vempala [15] in the  $\psi_1$  case. However,

in our approach  $t$  is going to be some fixed number which is bounded away from 0, thus optimizing over it cannot yield better bounds.

**Proposition 2.2.** *Suppose  $K$  is a convex body centered at the origin which is  $\psi_\alpha$  with constant  $b_\alpha$ . Then, for some universal constant  $c > 0$ ,*

$$\Delta_{KB}(K) \geq \frac{\exp(cb_\alpha^{-\alpha}n^{\alpha/2})}{2^n}.$$

We remark that Theorem 1.3 is a particular consequence of Proposition 2.2, as all random vectors with log-concave densities are  $\psi_1$  with the same universal constant.

*Proof of Proposition 2.2.* Let  $X$  and  $Y$  be independent random vectors, uniformly distributed on  $K$ . Since  $\Delta_{KB}(K)$  is affine invariant, we may assume without loss of generality that  $K$  is in isotropic position: this means that  $|K| = 1$ ,  $b(K) = 0$  as assumed already, and that  $\mathbb{E}(X \otimes X)$  is a multiple of the identity,  $\mathbb{E}(X \otimes X) = L_K^2 Id$  where  $L_K$  is called the isotropic constant of  $K$  (note that this is another well-defined affine invariant of  $K$ ). Equivalently, we ask that  $|K| = 1$  and  $X/L_K$  is isotropic as defined above.

We are now looking for a lower bound for  $\|f\|_\infty$  where  $f = \mathbb{1}_K * \mathbb{1}_K$  is the density function for the random vector  $X + Y$ . Instead, we shall work with  $\frac{X+Y}{2}$  so that both  $\frac{X+Y}{2}$  and  $X$  have the same support. The probability density function of  $\frac{X+Y}{2}$  is then  $g(x) = f(2x)2^n$ . There are many nice properties that  $\frac{X+Y}{2}$  inherits from  $X$ . In particular,  $\frac{X+Y}{2}$  has a centered log-concave density (the latter is a consequence of the Prékopa-Leindler inequality, see e.g. [2]). Moreover,

$$\begin{aligned} & \mathbb{E}_{X,Y} \left( \frac{X+Y}{2} \right) \otimes \left( \frac{X+Y}{2} \right) \\ &= \frac{1}{4} \mathbb{E}_{X,Y} (X \otimes X + X \otimes Y + Y \otimes X + Y \otimes Y) \\ &= \frac{1}{4} (L_K^2 I + 0 + 0 + L_K^2 I) \\ &= \frac{1}{2} L_K^2 I. \end{aligned}$$

Thus,  $\frac{X+Y}{2}$  is isotropic up to scaling. Finally,  $\frac{X+Y}{2}$  has more or less the same  $\psi_\alpha$  behavior as  $X$  (indeed, the above computations already show that

$$\left( \mathbb{E} \left| \left\langle \frac{X+Y}{2}, y \right\rangle \right|^2 \right)^{1/2} = \frac{1}{\sqrt{2}} L_K \|y\|_2 = \frac{1}{\sqrt{2}} \left( \mathbb{E} \left| \langle X, y \rangle \right|^2 \right)^{1/2}$$

for every  $y \in \mathbb{R}^n$ , hence a single application of Minkowski's inequality gives

$$\begin{aligned} \left( \mathbb{E} \left| \left\langle \frac{X+Y}{2}, y \right\rangle \right|^p \right)^{1/p} &\leq 2 \left( \mathbb{E} \left| \left\langle \frac{X}{2}, y \right\rangle \right|^p \right)^{1/p} = \left( \mathbb{E} \left| \langle X, y \rangle \right|^p \right)^{1/p} \\ &\leq b_\alpha p^{1/\alpha} \left( \mathbb{E} \left| \langle X, y \rangle \right|^2 \right)^{1/2} = \sqrt{2} b_\alpha p^{1/\alpha} \left( \mathbb{E} \left| \left\langle \frac{X+Y}{2}, y \right\rangle \right|^2 \right)^{1/2}, \end{aligned}$$

assuming  $X$  is  $\psi_\alpha$  with constant  $b_\alpha$ ). It is worth remarking however that, for our proof here, the fact that the distribution of  $\frac{X+Y}{2}$  is  $\psi_1$  suffices (and, as mentioned already, this is true for every log-concave distribution).

Observe now that for any  $r > 0$  we have

$$\|g\|_\infty \geq \frac{\int_{rL_K\sqrt{n}B_2^n \cap K} g(x) \, dx}{\int_{rL_K\sqrt{n}B_2^n \cap K} 1 \, dx} = \frac{\mathbb{P}(\|\frac{X+Y}{2}\|_2 \leq rL_K\sqrt{n})}{\mathbb{P}(\|X\|_2 \leq rL_K\sqrt{n})}.$$

Since  $\mathbb{E}_{X,Y} \|\frac{X+Y}{2}\|_2^2 = \frac{1}{2}nL_K^2$  and  $\mathbb{E}_X \|X\|_2^2 = nL_K^2$ , we know that the distributions of  $X$  and  $\frac{X+Y}{2}$  are concentrated within two different thin-shells. Thus, for  $\frac{1}{\sqrt{2}} < r < 1$ , we get that  $\mathbb{P}_{X,Y}(\|\frac{X+Y}{2}\|_2 \leq rL_K\sqrt{n})$  is almost 1 since the set considered includes the “good” thin-shell of  $\frac{X+Y}{2}$ . On the other hand,  $\mathbb{P}(\|X\|_2 \leq rL_K\sqrt{n})$  is almost 0 since the set considered excludes the corresponding thin-shell of  $X$ . To quantify this, we apply Theorem 2.1: for any isotropic  $\psi_\alpha$  log-concave vector  $Z$  the inequality in 2.1 is split into

$$\mathbb{P}(\|Z\|_2 \leq (1-t)\sqrt{n}) \leq C \exp(-c'b_\alpha^{-\alpha} \min(t^{2+\alpha}, t)\sqrt{n}) \quad \forall t \in [0, 1],$$

$$\mathbb{P}(\|Z\|_2 \geq (1+t)\sqrt{n}) \leq C \exp(-c'b_\alpha^{-\alpha} \min(t^{2+\alpha}, t)\sqrt{n}) \quad \forall t \geq 0.$$

Since we shall apply the first one with  $Z$  replaced by  $X/L_K$  and the second one with  $Z$  replaced by  $\frac{X+Y}{2} \cdot \frac{\sqrt{2}}{L_K}$ , we need  $1-t = \frac{1+t}{\sqrt{2}}$  and hence  $t = \frac{\sqrt{2}-1}{\sqrt{2}+1}$ . We thus obtain

$$\mathbb{P}\left(\|X\|_2 \leq \frac{2}{\sqrt{2}+1}L_K\sqrt{n}\right) \leq \exp(-c'b_\alpha^{-\alpha}n^{\alpha/2}),$$

and

$$\mathbb{P}\left(\left\|\frac{X+Y}{2}\right\|_2 \leq \frac{2}{\sqrt{2}+1}L_K\sqrt{n}\right) \geq 1 - \exp(-c'b_\alpha^{-\alpha}n^{\alpha/2}).$$

Therefore, we conclude that for some universal constant  $c > 0$

$$\|g\|_\infty \geq \exp(cb_\alpha^{-\alpha}n^{\alpha/2}),$$

and equivalently

$$\Delta_{KB}(K) = \frac{\|g\|_\infty}{2^n} \geq \frac{\exp(cb_\alpha^{-\alpha}n^{\alpha/2})}{2^n}.$$

□

### 3. A NEW BOUND FOR HADWIGER'S COVERING PROBLEM

This section is devoted to the proof of Theorems 1.1 and 1.2. To that end, we need some preliminaries.

Let  $\bar{N}(A, B) = \min\left\{N : \exists x_1, \dots, x_N \in A \text{ such that } A \subseteq \bigcup_{i=1}^N \{x_i + B\}\right\}$  be the covering number of  $A$  by translates of  $B$  that are centered in  $A$ . We shall need the following volume ratio bound.

**Lemma 3.1.** *Let  $A, B \subseteq \mathbb{R}^n$  be convex bodies. Suppose  $B$  contains the origin in its interior. Then*

$$\overline{N}(A, B) \leq 2^n \frac{|A + \frac{1}{2}(B \cap (-B))|}{|B \cap (-B)|}.$$

*Proof.* Recall that the separation number of  $A$  in  $B$  is defined as

$$M(A, B) = \max\{M : \exists x_1, \dots, x_M \in A \text{ such that } \forall i \neq j (x_i + B) \cap (x_j + B) = \emptyset\}.$$

It is an easy exercise (see e.g. [3]) to show that

$$M(A, B) \leq \frac{|A + B|}{|B|}.$$

Next, note that for any convex body  $T \subseteq \mathbb{R}^n$ , one has  $\overline{N}(A, T - T) \leq M(A, T)$ . Indeed, take a maximal  $T$ -separated set in  $A$ , that is a set of points  $x_1, \dots, x_M \in A$  such that for every point  $x \in A$  one has  $(x + T) \cap \bigcup_{i=1}^M \{x_i + T\} \neq \emptyset$ . This means that  $A \subseteq \bigcup_{i=1}^M \{x_i + T - T\}$  or, in other words, that  $\overline{N}(A, T - T) \leq M(A, T)$ . Since  $\overline{N}(A, B) \leq \overline{N}(A, B \cap (-B))$ , it follows that  $\overline{N}(A, B) \leq M(A, \frac{1}{2}(B \cap (-B)))$ , and hence

$$\overline{N}(A, B) \leq 2^n \frac{|A + \frac{1}{2}(B \cap (-B))|}{|B \cap (-B)|}.$$

□

Next, we recall the notion of fractional covering numbers, as defined in [3]. Remember that  $\mathbb{1}_A$  stands for the indicator function of a set  $A \subseteq \mathbb{R}^n$ . A sequence of pairs  $S = \{(x_i, \omega_i) : x_i \in \mathbb{R}^n, \omega_i \in \mathbb{R}^+\}_{i=1}^N$  of points and weights is said to be a fractional covering of a set  $K \subseteq \mathbb{R}^n$  by a set  $T \subseteq \mathbb{R}^n$  if for all  $x \in K$  we have  $\sum_{i=1}^N \omega_i \mathbb{1}_{x_i+T}(x) \geq 1$ . The total weight of the covering is denoted by  $\omega(S) = \sum_{i=1}^N \omega_i$ . The fractional covering number of  $K$  by  $T$  is defined to be the infimal total weight over all fractional coverings of  $K$  by  $T$  and is denoted by  $N_\omega(K, T)$ .

We shall also need the following volume ratio bound from [3]:

**Lemma 3.2** ([3, Proposition 2.9]). *Let  $K, T \subseteq \mathbb{R}^n$  be convex bodies. Then*

$$N_\omega(K, T) \leq \frac{|K - T|}{|T|}.$$

Finally, we shall need the following inequality that relates integral covering numbers and fractional covering numbers, and which was proved in [3], and improved in [22] through the removal of lower order terms.

$$(3.1) \quad N(K, T_1 + T_2) \leq N_\omega(K, T_1)(1 + \ln \overline{N}(K, T_2)).$$



*Proof of Theorem 1.1.* We can assume without loss of generality that  $b(K) = 0$ . By Lemma 3.2, for  $0 < \lambda < 1$  and any  $x \in \mathbb{R}^n$  we have

$$\begin{aligned} N_\omega(K, \lambda K) &\leq N_\omega(K, \lambda(K \cap (x - K))) \leq \frac{|K - \lambda(K \cap (x - K))|}{|\lambda(K \cap (x - K))|} \\ &\leq \left(\frac{1 + \lambda}{\lambda}\right)^n \frac{|K|}{|K \cap (x - K)|}. \end{aligned}$$

By applying Theorem 1.3 with the point  $x$  which maximizes the above volume ratio, we get

$$N_\omega(K, \lambda K) \leq \left(\frac{1 + \lambda}{\lambda}\right)^n 2^n e^{-c\sqrt{n}}.$$

Using (3.1) with  $T_1 = \alpha\lambda K, T_2 = (1 - \alpha)\lambda K$  for some  $\alpha \in (0, 1)$ , we obtain

$$N(K, \lambda K) \leq \left(\frac{1 + \alpha\lambda}{\alpha\lambda}\right)^n 2^n e^{-c\sqrt{n}} (1 + \ln \bar{N}(K, (1 - \alpha)\lambda K)).$$

Using Lemma 3.1 and taking the limit  $\lambda \nearrow 1$ , we get

$$\begin{aligned} N(K, \text{int}K) &\leq \left(\frac{1 + \alpha}{\alpha}\right)^n 2^n e^{-c\sqrt{n}} \left(1 + \ln \left(2^n \frac{|K + \frac{1}{2}(1 - \alpha)(K \cap (-K))|}{|(1 - \alpha)(K \cap (-K))|}\right)\right) \\ &\leq \left(\frac{1 + \alpha}{\alpha}\right)^n 2^n e^{-c\sqrt{n}} \left(1 + \ln \left(\left(\frac{4}{1 - \alpha}\right)^n \frac{|K|}{|K \cap (-K)|}\right)\right). \end{aligned}$$

Since  $K$  is centered at the origin, (1.2) (or its improvement in Theorem 1.3, which however cannot essentially affect the final estimate here) implies that

$$\begin{aligned} N(K, \text{int}K) &\leq \left(\frac{1 + \alpha}{\alpha}\right)^n 2^n e^{-c\sqrt{n}} \left(1 + \ln \left(\left(\frac{4}{1 - \alpha}\right)^n 2^n\right)\right) \\ &\leq \left(\frac{1 + \alpha}{\alpha}\right)^n 2^n e^{-c\sqrt{n}} \left(1 + n \ln \left(\frac{8}{1 - \alpha}\right)\right). \end{aligned}$$

Plugging in  $\alpha = 1 - 1/n$  yields that, for some universal constants  $c_1, c_2 > 0$ , we have

$$N(K, \text{int}K) \leq c_1 4^n e^{-c_2\sqrt{n}}.$$

□

The proof of Theorem 1.2 is the same as that of Theorem 1.1, except that one uses Corollary 1.4 instead of Theorem 1.3.

#### 4. POSITIVE MODULUS OF CONVEXITY

Recall that the modulus of convexity of a centered convex body  $K \subseteq \mathbb{R}^n$  is defined by

$$\delta_K(\varepsilon) = \inf \left\{ 1 - \left\| \frac{x + y}{2} \right\|_K : \|x\|_K, \|y\|_K \leq 1, \|x - y\|_K \geq \varepsilon \right\}.$$

A result of Arias-De-Reyna, Ball, and Villa [1], which was generalized by Gluskin and Milman [9], tells us that if  $K \subseteq \mathbb{R}^n$  is a convex body such that  $0 \in \text{int}K$  and

$|K| = 1$  then for all  $0 < \varepsilon' < 1$  one has

$$(4.1) \quad \left| \left\{ (x, y) \in K \times K : \|x - y\|_K \leq \sqrt{2}(1 - \varepsilon') \right\} \right| \leq e^{-\varepsilon'^2 n/2}.$$

We use this result to prove Theorem 1.5:

*Proof of first part of Theorem 1.5.* Without loss of generality, we assume that  $|K| = 1$ . Let  $X$  and  $Y$  be independent random vectors, uniformly distributed on  $K$ . Let  $f(x) = |K \cap (x - K)|$  and recall that the density of  $\frac{X+Y}{2}$  is  $g(x) = 2^n f(2x)$ .

Since, by assumption,  $\delta_K(\varepsilon) \geq r$ , the set  $\theta = \{(x, y) \in K \times K : \|x - y\|_K \geq \varepsilon\}$  satisfies that

$$\theta \subseteq \left\{ (x, y) \in K \times K : \frac{x + y}{2} \in (1 - r)K \right\}.$$

By (4.1), we have that  $|\theta| \geq 1 - e^{-\frac{(\sqrt{2}-\varepsilon)^2 n}{4}}$  and hence

$$\begin{aligned} \mathbb{P}\left(\frac{X + Y}{2} \in (1 - r)K\right) &= \iint_{\{(x, y) \in K \times K : \frac{x+y}{2} \in (1-r)K\}} dx dy \\ &\geq \iint_{\theta} dx dy \geq 1 - e^{-\frac{(\sqrt{2}-\varepsilon)^2 n}{4}}. \end{aligned}$$

Therefore, it follows that

$$\begin{aligned} \|g\|_{\infty} &\geq \frac{\int_{(1-r)K} g(x) dx}{\int_{(1-r)K} dx} = \frac{\mathbb{P}\left(\frac{X+Y}{2} \in (1-r)K\right)}{\mathbb{P}(X \in (1-r)K)} \\ &\geq \left(\frac{1}{1-r}\right)^n \left(1 - e^{-\frac{(\sqrt{2}-\varepsilon)^2 n}{4}}\right). \end{aligned}$$

□

Repeating the proof of Theorem 1.1 but now using Theorem 1.5, Theorem 1.6 follows.

## 5. BOUNDING THE CONVOLUTION SQUARE AT THE BARYCENTER

This section is devoted to the proof of Proposition 5.3 below (which will give the full proofs of Theorem 1.3 and Corollary 1.4) as well as completing that of Theorem 1.5 (the arguments will be very similar, just different applications of the same method). We recall that, for a random vector  $X$  in  $\mathbb{R}^n$  with density  $f$ , we define its entropy as

$$\text{Ent}[X] = - \int_{\mathbb{R}^n} f \ln f.$$

The conclusions of the following standard lemma are simple consequences of Jensen's inequality.

**Lemma 5.1.** *For any measurable function  $h : \mathbb{R}^n \rightarrow [0, +\infty)$  which is positive on the support of  $f$  we have*

$$(5.1) \quad \text{Ent}[X] \leq - \int_{\mathbb{R}^n} f \ln h + \ln \left( \int_{\mathbb{R}^n} h \right),$$

*assuming all the quantities are finite. Moreover, if  $X$  has a log-concave density, then*

$$(5.2) \quad \text{Ent}[X] = \mathbb{E}[-\ln f(X)] \geq -\ln f(\mathbb{E}X).$$

*Proof.* To prove (5.1), we write

$$\text{Ent}[X] + \int_{\mathbb{R}^n} f \ln h = \int_{\mathbb{R}^n} f \ln \frac{h}{f} \leq \ln \left( \int_{\mathbb{R}^n} h \right),$$

with the inequality following by Jensen's inequality. As for (5.2), we note that, if  $f$  is assumed log-concave,  $-\log f$  will be a convex function on  $\mathbb{R}^n$ , which allows to apply Jensen's inequality again.  $\square$

*Remark 5.2.* We will apply Lemma 5.1 as follows. If  $K \subset \mathbb{R}^n$  is a centered convex body, and  $X, Y$  are independent random vectors uniformly distributed on  $K$ , then the density  $f$  of  $X$  is given by  $f(x) = \frac{1}{|K|} \mathbb{1}_K$ , while the density  $g$  of  $X + Y$  by  $g(x) = \frac{1}{|K|^2} (\mathbb{1}_K * \mathbb{1}_K)(x) = \frac{1}{|K|^2} |K \cap (x - K)|$  (recall that  $X + Y$  has a centered log-concave density, which is not hard to check using this identity). These show that  $\text{Ent}[X] = \log |K|$ , while, by (5.2),

$$-\ln \left( \frac{|K \cap (-K)|}{|K|^2} \right) = -\ln g(0) \leq \text{Ent}[X + Y].$$

Therefore,

$$(5.3) \quad -\ln \left( \frac{|K \cap (-K)|}{|K|} \right) = -\ln \left( \frac{|K \cap (-K)|}{|K|^2} \right) - \ln |K| \leq \text{Ent}[X + Y] - \text{Ent}[X],$$

which we can combine with (5.1), applied for the vector  $X + Y$ , to obtain

$$(5.4) \quad -\ln \left( \frac{|K \cap (-K)|}{|K|} \right) \leq \mathbb{E}[-\ln h(X + Y)] + \ln \left( \int_{\mathbb{R}^n} h \right) - \text{Ent}[X]$$

for any integrable function  $h : \mathbb{R}^n \rightarrow [0, +\infty)$  which is positive on  $2K$  (note that the first term on the right hand side depends only on values of  $h$  on  $2K$ , whereas the second term can only get smaller or stay the same when  $h$  is restricted to  $2K$ ; in other words, replacing  $h$  with  $h \mathbb{1}_{2K}$  might only improve the right hand side).

Observe that, by choosing  $h$  constant on  $2K$  (and zero otherwise), one can recover (1.2). In the remainder of this section, we will choose different  $h$  in order to establish the improvements of (1.2) claimed earlier.

**Proposition 5.3.** *Suppose  $K$  is a convex body centered at the origin which is  $\psi_\alpha$  with constant  $b_\alpha$ . Then, for some universal constant  $c > 0$ ,*

$$\frac{|K \cap (-K)|}{|K|} \geq \frac{\exp(cb_\alpha^{-\alpha}n^{\alpha/2})}{2^n}.$$

*Proof.* We begin by observing that both sides of (5.3) are invariant under invertible linear transformations of  $K$ , therefore we can assume without loss of generality that  $K$  is in isotropic position. We then apply (5.4) with  $h(x) := \exp(-\lambda\|x\|_2^2)\mathbb{1}_{2K}$  for some constant  $\lambda$  to be specified later. The right hand side becomes

$$\begin{aligned} \mathbb{E}[\lambda\|X + Y\|_2^2] + \ln \int_{2K} \exp(-\lambda\|x\|_2^2) dx - \ln 1 &= 2\mathbb{E}[\lambda\|X\|_2^2] + \ln \int_{2K} \exp(-\lambda\|x\|_2^2) dx \\ (5.5) \qquad \qquad \qquad &= 2\lambda nL_K^2 + n \ln 2 + \ln \int_K \exp(-4\lambda\|x\|_2^2) dx. \end{aligned}$$

To estimate the last integral, we employ again the thin-shell estimates from Theorem 2.1, which imply that for  $A_t := \{x \in K : \|x\|_2 \leq (1-t)\sqrt{n}L_K\}$ , one has

$$|A_t| \leq C \exp(-c'b_\alpha^{-\alpha}t^{2+\alpha}n^{\alpha/2})$$

for all  $t \in [0, 1]$ . We can thus break the integral into two as follows:

$$\begin{aligned} \int_K \exp(-4\lambda\|x\|_2^2) dx &= \int_{A_t} \exp(-4\lambda\|x\|_2^2) dx + \int_{K \setminus A_t} \exp(-4\lambda\|x\|_2^2) dx \\ &\leq C \exp(-c'b_\alpha^{-\alpha}t^{2+\alpha}n^{\alpha/2}) + \exp(-4\lambda(1-t)^2nL_K^2). \end{aligned}$$

We now set  $t = 1 - \frac{2}{\sqrt{5}}$  say, and then we choose our  $\lambda$  so that

$$c'b_\alpha^{-\alpha}t^{2+\alpha}n^{\alpha/2} = 4\lambda(1-t)^2nL_K^2.$$

It follows that  $\lambda$  is of the order of  $b_\alpha^{-\alpha}n^{\alpha/2-1}L_K^{-2}$ . Combining these estimates with (5.4) and (5.5), we obtain

$$\begin{aligned} -\ln\left(\frac{|K \cap (-K)|}{|K|}\right) &\leq 2\lambda nL_K^2 + n \ln 2 + \ln(C+1) - \frac{16}{5}\lambda nL_K^2 \\ &= n \ln 2 + \ln(C+1) - \frac{6}{5}\lambda nL_K^2 \\ &= n \ln 2 + \ln(C+1) - c''b_\alpha^{-\alpha}n^{\alpha/2} \end{aligned}$$

for some absolute constant  $c''$  (which we can compute explicitly by the above relations). Exponentiating, we complete the proof.  $\square$

*Proof of second part of Theorem 1.5.* This time we only assume for simplicity that  $|K| = 1$ , and we apply (5.4) with  $h(x) := \exp(-\lambda\|x\|_K)$  for some constant  $\lambda$  to be specified later. We immediately get

$$\begin{aligned} -\ln\left(\frac{|K \cap (-K)|}{|K|}\right) &\leq \mathbb{E}[\lambda\|X + Y\|_K] + \ln \int_{\mathbb{R}^n} \exp(-\lambda\|x\|_K) dx \\ &= \lambda\mathbb{E}[\|X + Y\|_K] + \ln(\lambda^{-n}n!|K|) = \lambda\mathbb{E}[\|X + Y\|_K] - n \ln \lambda + \ln(n!). \end{aligned}$$

Optimizing over  $\lambda$  yields

$$(5.6) \quad -\log\left(\frac{|K \cap (-K)|}{|K|}\right) \leq n \ln \mathbb{E}[\|X + Y\|_K] + \ln \frac{n!e^n}{n^n}.$$

Given that  $n! \leq en^{n+1/2}e^{-n}$ , the last term is upper-bounded by  $\log(e\sqrt{n})$ , so the final estimate will depend on how well we can bound  $\mathbb{E}[\|X + Y\|_K]$ . We will use again the concentration result of Arias-De-Reyna, Ball, and Villa. Note that by the triangle inequality  $\|X + Y\|_K \leq 2$ , and therefore, by the definition of the modulus of convexity, we have for any  $\varepsilon \in (0, 2)$ ,

$$\begin{aligned} \mathbb{E}[\|X + Y\|_K] &= \mathbb{E}[\|X + Y\|_K \mathbb{1}_{\|X - Y\|_K \leq \varepsilon}] + \mathbb{E}[\|X + Y\|_K \mathbb{1}_{\|X - Y\|_K > \varepsilon}] \\ &\leq 2\mathbb{P}(\|X - Y\|_K \leq \varepsilon) + 2(1 - \delta_K(\varepsilon))\mathbb{P}(\|X - Y\|_K > \varepsilon) \\ &= 2[1 - \delta_K(\varepsilon)\mathbb{P}(\|X - Y\|_K > \varepsilon)]. \end{aligned}$$

Applying this now with some  $\varepsilon \in (0, \sqrt{2})$  for which  $\delta_K(\varepsilon) \geq r$ , and recalling (4.1), we obtain

$$\mathbb{E}[\|X + Y\|_K] \leq 2 \left[ 1 - \delta_K(\varepsilon) \left( 1 - \exp\left(-\frac{(\sqrt{2}-\varepsilon)^2 n}{4}\right) \right) \right] \leq 2 \left[ 1 - r \left( 1 - \exp\left(-\frac{(\sqrt{2}-\varepsilon)^2 n}{4}\right) \right) \right],$$

which we can plug into (5.6) to complete the proof.  $\square$

## 6. CONCLUDING REMARKS

We conclude this note with some remarks, questions and conjectures.

**Conjecture 6.1.** *There exists a universal constant  $c > 0$  such that for every centered convex body  $K \subseteq \mathbb{R}^n$  and some  $0 < r < 1$  one has*

$$\frac{\mathbb{P}\left(\frac{X+Y}{2} \in rK\right)}{\mathbb{P}(X \in rK)} \geq (1+c)^n,$$

where  $X$  and  $Y$  are independent random vectors, uniformly distributed on  $K$ .

We remark that the above conjecture implies an exponentially better upper bound for Hadwiger's covering problem. Moreover, the conjecture seems interesting in its

own right, even for centrally-symmetric convex bodies, although in this case it will have no meaningful implication towards Hadwiger's problem.

Note that this conjecture attempts in a way to quantify the intuition that the convolution of a uniform distribution with itself looks already more like a "bell curve" than like the flat distribution it originates from. In other words, we expect that the sum of two independent copies of a uniform random vector  $X$  concentrates much better around the origin. Another question that would capture this if answered in the affirmative is the following. Recall that a simple application of Fubini's theorem and the homogeneity of the Lebesgue volume give

$$\mathbb{E}\|X\|_K = 1 - \frac{1}{n+1},$$

thus, simply by the triangle inequality, we have  $\mathbb{E}\|X + Y\|_K \leq 2 - \frac{2}{n+1}$  whenever  $X, Y$  are as above. Note however that, because of independence, we might expect the following.

**Question 6.2.** What is the behavior of  $\sup_{b(K)=0} \mathbb{E}\|X + Y\|_K$ ? Is it better than what the bound following by the triangle inequality suggests (either because it is upper-bounded by some constant smaller than 2 or because it converges to 2 slower)?

In the previous section we proved that the linearly invariant quantity  $\mathbb{E}\|X + Y\|_K$  is indeed upper-bounded by a constant smaller than 2 for convex bodies with a positive modulus of convexity. One might be tempted to think that, even though this proof would obviously not work for general bodies, maybe such an upper bound still holds. However, this is not true, and surprisingly perhaps one example showing this is the  $n$ -cube  $Q_n \equiv [-1, 1]^n$ : using the fact that its uniform distribution is a product measure, we can compute as follows.

$$\begin{aligned} \frac{1}{|Q_n|^2} \int_{Q_n} \int_{Q_n} \|x + y\|_{Q_n} dx dy &= \int_0^2 \mathbb{P}(\|X + Y\|_{Q_n} \geq t) dt \\ &= \int_0^2 \mathbb{P}\left(\max_i |X_i + Y_i| \geq t\right) dt \\ &= \int_0^2 \left(1 - \mathbb{P}\left(\max_i |X_i + Y_i| < t\right)\right) dt \\ &= 2 - \int_0^2 [\mathbb{P}(|X_1 + Y_1| < t)]^n dt. \end{aligned}$$

To compute the last integral, we first find the convolution square  $g$  of the indicator function of the interval  $[-1, 1]$ :  $g$  is supported on  $[-2, 2]$  and for every  $s \in [-2, 2]$  we have  $g(s) = \frac{2-|s|}{4}$ . Therefore, for every  $t \in [0, 2]$ ,

$$\mathbb{P}(|X_1 + Y_1| < t) = \int_{-t}^t g(s) ds = \frac{1}{4}t(4 - t),$$

and returning to the integrals above we get

$$\begin{aligned} \frac{1}{|Q_n|^2} \int_{Q_n} \int_{Q_n} \|x + y\|_{Q_n} dx dy &= 2 - \frac{1}{4^n} \int_0^2 t^n (4 - t)^n dt \\ &= 2 - \frac{1}{2 \cdot 4^n} \int_0^4 t^n (4 - t)^n dt \\ (t = 4s) \qquad &= 2 - 2 \cdot 4^n \frac{\Gamma(n+1)\Gamma(n+1)}{\Gamma(2n+2)} \\ &= 2 - \frac{2}{2n+1} 4^n \binom{2n}{n}^{-1}. \end{aligned}$$

This finally gives

$$2 - \frac{\sqrt{2\pi}}{\sqrt{2n+1}} \leq \frac{1}{|Q_n|^2} \int_{Q_n} \int_{Q_n} \|x + y\|_{Q_n} dx dy \leq 2 - \frac{\sqrt{\pi}}{\sqrt{2n+1}}$$

by standard numerical estimates for  $\binom{2n}{n}$  (see e.g. [28, Lemma 4.2.4]). We could make the following conjecture.

**Conjecture 6.3.** For the convex body/-ies  $K_0$  at which  $\sup_{\mathfrak{b}(K)=0} \mathbb{E}\|X+Y\|_K$  is attained the behavior of the relevant quantity is similar to that for the  $n$ -cube.

We could also make the stronger conjecture that  $\sup_{\mathfrak{b}(K)=0} \mathbb{E}\|X+Y\|_K$  is attained at the cube. Note that, if either form of the conjecture has an affirmative answer, then the approach discussed in the previous section for uniformly convex bodies could be extended to give an alternative proof of Theorem 1.3. The conjecture of course seems interesting in its own right, again giving some evidence towards the intuition discussed above.

The quantity  $\text{Ent}[X+Y] - \text{Ent}[X]$  which appears on the right hand side of (5.3) has been studied in the context of reverse entropy power inequalities for convex measures, a natural generalization of log-concave measures (see [5] and [19]). The upper bounds obtained there (when specialized to the log-concave case) as well as our improved bounds are perhaps far from optimal. To the best of our knowledge, a sharp upper bound is not known even in dimension one. We believe the extremizer would be a one-sided exponential distribution.

*Acknowledgement.* We thank the Mathematical Sciences Research Institute in Berkeley, California, where part of this work was done during the Fall 2017 semester; we acknowledge the support of the institute and of the National Science Foundation

under Grant No. 1440140. We also thank Shiri Artstein-Avidan and Bo'az Klartag for useful discussions.

## REFERENCES

- [1] J. Arias-de Reyna, K. Ball, and R. Villa, *Concentration of the distance in finite dimensional normed spaces*, *Mathematika* **45** (1998), no. 2, 245–252.
- [2] S. Artstein-Avidan, A. Giannopoulos, and V. D. Milman, *Asymptotic geometric analysis. Part I*, *Mathematical Surveys and Monographs*, vol. 202, American Mathematical Society, Providence, RI, 2015.
- [3] S. Artstein-Avidan and B. A. Slomka, *On weighted covering numbers and the Levi-Hadwiger conjecture*, *Israel J. Math.* **209** (2015), no. 1, 125–155.
- [4] K. Bezdek and M. A. Khan, *The geometry of homothetic covering and illumination*, *Discrete Geometry and Symmetry (Cham)* (M. D. E. Conder, A. Deza, and A. I. Weiss, eds.), Springer International Publishing, 2018, pp. 1–30.
- [5] S. G. Bobkov and M. M. Madiman, *On the problem of reversibility of the entropy power inequality*, *Limit theorems in probability, statistics and number theory*, *Springer Proc. Math. Stat.*, vol. 42, Springer, Heidelberg, 2013, pp. 61–74.
- [6] V. Boltyanski, *A problem on the illumination of the boundary of a convex body (in Russian)*, *Izv. Mold. Fil. Akad. Nauk SSSR* (1960), no. 10, 77–84.
- [7] P. Brass, W. Moser, and J. Pach, *Research problems in discrete geometry*, Springer, New York, 2005.
- [8] P. Erdős and C. A. Rogers, *The star number of coverings of space with convex bodies*, *Acta Arith.* **9** (1964), 41–45.
- [9] E. Gluskin and V. Milman, *Geometric probability and random cotype 2*, pp. 123–138, Springer Berlin Heidelberg, Berlin, Heidelberg, 2004.
- [10] I. Gohberg and A. Markus, *A problem on covering of convex figures by similar figures (in Russian)*, *Izv. Mold. Fil. Akad. Nauk SSSR* **10** (1960), no. 76, 87–90.
- [11] B. Grünbaum, *Measures of symmetry for convex sets*, *Proc. Sympos. Pure Math.*, Vol. VII, Amer. Math. Soc., Providence, R.I., 1963, pp. 233–270.
- [12] O. Guédon and E. Milman, *Interpolating Thin-Shell and Sharp Large-Deviation Estimates for Isotropic Log-Concave Measures*, *Geometric and Functional Analysis* **21** (2011), no. 5, 1043–1068.
- [13] H. Hadwiger, *Ungelöstes Probleme Nr. 20*, *Elem. Math.* **12** (1957), no. 6, 121.
- [14] ———, *Ungelöstes Probleme Nr. 38*, *Elem. Math.* **15** (1960), 130–131.
- [15] Y. T. Lee and S. S. Vempala, *Stochastic localization + stieljes barrier = tight bound for log-sobolev*, *Proceedings of the 50th Annual ACM SIGACT Symposium on Theory of Computing (New York, NY, USA)*, *STOC 2018*, ACM, 2018, pp. 1122–1129.
- [16] F. W. Levi, *Überdeckung eines Eibereiches durch Parallelverschiebung seines offenen Kerns*, *Arch. Math. (Basel)* **6** (1955), 369–370.
- [17] G. Livshyts and K. Tikhomirov, *Randomized coverings of a convex body with its homothetic copies, and illumination*, *ArXiv e-prints* (2016), arXiv:1606.08876.
- [18] ———, *Cube is a strict local maximizer for the illumination number*, *ArXiv e-prints* (2017), arXiv:1710.05070.
- [19] M. Madiman and I. Kontoyiannis, *Entropy bounds on abelian groups and the Ruzsa divergence*, *IEEE Trans. Inform. Theory* **64** (2018), no. 1, 77–92.



- [20] V. D. Milman and A. Pajor, *Entropy and Asymptotic Geometry of Non-Symmetric Convex Bodies*, *Advances in Mathematics* **152** (2000), no. 2, 314–335.
- [21] M. Naszódi, *Fractional illumination of convex bodies*, *Contributions to Discrete Mathematics* **4** (2009), no. 2, 83–88.
- [22] M. Naszódi, *On some covering problems in geometry*, *Proc. Amer. Math. Soc.* **144** (2016), no. 8, 3555–3562.
- [23] M. Naszódi, *Flavors of translative coverings*, pp. 335–358, Springer Berlin Heidelberg, Berlin, Heidelberg, 2018.
- [24] A. Prymak and V. Shepelska, *On illumination of the boundary of a convex body in  $\mathbb{E}^n$ ,  $n = 4, 5, 6$* , ArXiv e-prints (2018), arXiv:1811.08962.
- [25] C. A. Rogers, *A note on coverings*, *Mathematika* **4** (1957), no. 1, 1–6.
- [26] C. A. Rogers and G. C. Shephard, *The difference body of a convex body*, *Arch. Math. (Basel)* **8** (1957), 220–233.
- [27] C. A. Rogers and C. Zong, *Covering convex bodies by translates of convex bodies*, *Mathematika* **44** (1997), no. 1, 215–218.
- [28] S. Taschuk, *The harmonic mean measure of symmetry for convex bodies*, *Adv. Geom.* **15** (2015), no. 1, 109–120.
- [29] K. Tikhomirov, *Illumination of convex bodies with many symmetries*, ArXiv e-prints (2016), arXiv:1606.08976.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, 2074 EAST HALL, 530 CHURCH ST., ANN ARBOR, MI 48109-1043, USA

*E-mail address:* sthhan@umich.edu

DEPARTMENT OF MATHEMATICS, WEIZMANN INSTITUTE OF SCIENCE, REHOVOT 76100, ISRAEL

*E-mail address:* boazslomka@gmail.com

DEPARTMENT OF MATHEMATICAL SCIENCES, CARNEGIE MELLON UNIVERSITY, WEAN HALL 6113, PITTSBURGH, PA 15213, USA

*E-mail address:* ttkocz@math.cmu.edu

DEPARTMENT OF MATHEMATICAL AND STATISTICAL SCIENCES, UNIVERSITY OF ALBERTA, CAB 632, EDMONTON, AB, CANADA T6G 2G1

*E-mail address:* vritsiou@ualberta.ca